

## Convenient Synthesis, Separation, and X-ray Crystal Structure Determinations of 1(e),3(e),5(e)-Trimethylcycloborazane and 1(e),3(e),5(a)-Trimethylcycloborazane

Chaitanya K. Narula, Jerzy Fr. Janik, Eileen N. Duesler, Robert T. Paine,\* and Riley Schaeffer


Received March 4, 1986

1,3,5-Trimethylcycloborazane  $[\text{CH}_3\text{N}(\text{H})\text{BH}_2]_3$  is obtained as a mixture of eee and eea isomers in good yield from the reaction of  $\text{BH}_3\cdot\text{THF}$  and methylamine. The isomers are separated by fractional crystallization from benzene and characterized by mass, infrared, and  $^1\text{H}$ ,  $^{13}\text{C}$ , and  $^{11}\text{B}$  NMR spectroscopies. The eee isomer crystallizes in the monoclinic space group  $P2_1/c$ , with  $a = 8.514$  (2) Å,  $b = 13.577$  (4) Å,  $c = 8.927$  (2) Å,  $\beta = 112.09$  (2)°,  $Z = 4$ ,  $V = 956.1$  (4) Å<sup>3</sup>, and  $\rho = 0.89$  g cm<sup>-3</sup>. Least-squares refinement gave  $R_F = 8.3\%$  and  $R_{wF} = 6.1\%$ . The compound displays a six-membered ring structure with alternating boron and nitrogen atoms in a chair conformation. The methyl groups on the nitrogen atoms are all in equatorial positions. The average B-N bond distance is 1.565 Å. The eea isomer crystallizes in the monoclinic space group  $Cc$ , with  $a = 9.481$  (3) Å,  $b = 12.477$  (4) Å,  $c = 8.329$  (2) Å,  $\beta = 114.05$  (2)°,  $Z = 4$ ,  $V = 900.0$  (5) Å<sup>3</sup>, and  $\rho = 0.95$  g cm<sup>-3</sup>. Least-squares refinement gave  $R_F = 4.8\%$  and  $R_{wF} = 4.5\%$ . This isomer also has a six-membered ring structure with alternating boron and nitrogen atoms in a chair conformation. Two methyl groups are found in equatorial positions, and one methyl group is found in an axial position. The average B-N distance is 1.578 Å.


### Introduction

There is currently a great deal of interest in the utilization of simple boron-nitrogen compounds as precursors to polymers and solid-state materials. On the basis of tantalizing observations in the literature,<sup>1,2</sup> the relatively stable cycloborazanes (parent molecule  $\text{B}_3\text{H}_6\text{N}_3\text{H}_6$ ) appear to be one family of compounds with potential as solid-state-material precursors. With increased interest in cycloborazanes has come renewed attention to the development of routine syntheses for these molecules. In this regard, improved syntheses and characterization of 1,3,5-trimethylcycloborazane,  $[\text{CH}_3\text{N}(\text{H})\text{BH}_2]_3$  (**1**), are of interest to us.



Bissot and Parry<sup>3</sup> reported the first synthesis and characterization of **1** from the gentle thermolysis of the amine borane adduct  $\text{CH}_3\text{NH}_2\cdot\text{BH}_3$  at 100° (eq 1). Subsequently, Gaines and



Schaeffer<sup>4</sup> observed that **1** could be prepared in two isomeric forms from 1,3,5-trimethylborazane according to eq 2. More simply,



they observed that **1** is formed directly from the reaction of methylamine hydrochloride and sodium borohydride as described in eq 3. Brown and co-workers<sup>5</sup> also observed (eq 4) that thermal



decomposition of the appropriate amine boranes, produced  $[\text{RN}(\text{H})\text{BH}_2]_3$  with  $\text{R} = \text{CH}_3$ ,  $\text{C}_2\text{H}_5$ ,  $n\text{-C}_3\text{H}_7$ ,  $i\text{-C}_3\text{H}_7$ ,  $n\text{-C}_4\text{H}_9$ , and  $t\text{-C}_4\text{H}_9$ . Obviously there are several suitable laboratory routes to *N*-alkyl-substituted cycloborazanes. Reactions 1 and 2, in particular, provide good overall yields of the mixed isomers of **1**, 88% and 70%, respectively. However, the first reaction suffers from the disadvantage of utilizing  $\text{B}_2\text{H}_6$  as a starting material while the second reaction produces  $\text{B}_2\text{H}_6$  as a byproduct. The pyrophoric character of this gaseous reagent tends to reduce the attractiveness

of these reactions in process conditions. Reaction 3 is particularly convenient from a reactant standpoint and it circumvents the appearance of diborane; however, the yield of **1** is usually reduced (~50%) and is more susceptible to variations in experimental conditions.<sup>6</sup> We report here a simplified, convenient synthesis of **1** from commercially available and easily handled reagents. In addition, a complete set of characterization data and the single-crystal X-ray diffraction structure determinations for the two isomers are described.

### Experimental Section

**General Information.** All experimental manipulations were performed under rigorously anhydrous conditions in a high-vacuum system or in Schlenk equipment. Monoglyme was successively distilled from NaH,  $\text{LiAlH}_4$ , and sodium benzophenone ketyl. Commercial borane-tetrahydrofuran solutions (Aldrich) were used after checking their purity by  $^{11}\text{B}$  NMR. Methylamine was dried over KOH.

The  $^1\text{H}$ ,  $^{13}\text{C}$ , and  $^{11}\text{B}$  NMR spectra were recorded on Varian FT-80A or Nicolet 360-MHz spectrometers and referenced with  $(\text{CH}_3)_4\text{Si}$  ( $^1\text{H}$ ,  $^{13}\text{C}$ ) and  $\text{BF}_3\cdot\text{Et}_2\text{O}$  ( $^{11}\text{B}$ ). Infrared spectra were obtained on a Nicolet 6000 FTIR spectrometer from KBr pellets. Mass spectra were obtained on a Finnegan GC/MS system using a solids inlet probe.

**Preparation of  $[\text{CH}_3\text{N}(\text{H})\text{BH}_2]_3$  (**1**).** Borane-tetrahydrofuran complex (100 mL, 1 M solution; 100 mmol) was syringed into a dry, nitrogen-filled 250-mL Schlenk vessel containing a stir bar. The flask was frozen and degassed and dry methylamine (3.1 g, 100 mmol) condensed onto the solution at -78 °C. The flask was backfilled with nitrogen, and the mixture was slowly warmed to room temperature and then stirred at 25 °C for 1 h. The volatiles were removed in vacuo leaving a white, solid sample of  $\text{CH}_3\text{NH}_2\cdot\text{BH}_3$  (yield 4.33 g, 98%;  $^{11}\text{B}$  NMR,  $\delta$  -18.4). The resulting solid was dissolved in 15 mL of dry monoglyme, and the flask was outfitted with a condenser and nitrogen bubbler. The solution was refluxed under nitrogen until gas evolution ceased (~18 h), and the volatiles were removed in vacuo. The residue was washed with water, dried in air, and then sublimed (0.001 Torr, 70 °C). The sublimate was a white crystalline solid (yield 2.83 g, 69%), which contained a mixture of isomers I and II. The isomers were separated by dissolving the mixture in a minimum of hot benzene. Slow cooling of this solution deposited crystals of isomer I. When no evidence for isomer I was found in the solution by infrared analysis the crystals of isomer I were collected by filtration and washed with a minimum of cold benzene. The filtrate was slowly evaporated and crystalline isomer II was obtained [yields: isomer I 50%, isomer II 19% based upon  $\text{BH}_3\cdot\text{THF}$ ]. Analytical data melting points, sublimation temperatures, and solubility data for the isomers are in agreement with literature data.<sup>4</sup> Infrared spectra ( $\text{cm}^{-1}$ ) (KBr pellet): isomer I 3224 (s,  $\nu_{\text{NH}}$ ), 2995 (m,  $\nu_{\text{CH}}$ ) 2950 (w), 2351 (s,  $\nu_{\text{BH}}$ ), 2312 (w), 1460 (w), 1415 (w), 1367 (m), 1270 (m), 1193 (s), 1174 (s), 1151 (w), 1024 (w), 964 (w), 914 (w); isomer II 3276 (m,  $\nu_{\text{NH}}$ ), 3241 (m,  $\nu_{\text{NH}}$ ), 2996 (m,  $\nu_{\text{CH}}$ ), 2950 (w), 2890 (w), 2361 (s,  $\nu_{\text{BH}}$ ), 2342 (m,  $\nu_{\text{BH}}$ ), 1458 (m), 1417 (w), 1385 (m), 1365 (m), 1266 (w), 1233 (w), 1202 (s), 1177 (s), 1133 (w), 1065 (w), 1047 (w), 1021 (w), 984 (w), 962 (w), 925 (w), 908 (w). Mass spectra (70 eV) (unstripped): isomer I,  $m/e$  129 [ $\text{M}^+$ ]

(1) Wiberg, E. *Naturwiss* 1948 35, 212.

(2) Brown, M. P.; Heseltine, R. W. *J. Inorg. Nucl. Chem.* 1967, 29, 1197.

(3) Bissot, T. C.; Parry, R. W. *J. Am. Chem. Soc.* 1955, 77, 3481.

(4) Gaines, D. F.; Schaeffer, R. *J. Am. Chem. Soc.* 1963, 85, 395.

(5) Brown, M. P.; Heseltine, R. W.; Sutcliffe, L. H. *J. Chem. Soc. A* 1968, 612.

(6) Janik, J. Fr.; Narula, C. K., unpublished observations.

**Table I.** Experimental Data for the X-ray Diffraction Study of 1(e),3(e),5(e)-Trimethylcycloborazane (Isomer I) and 1(e),3(e),5(e)-Trimethylcycloborazane (Isomer II):  $\text{B}_3\text{N}_3\text{C}_3\text{H}_{18}$ 

	isomer I	isomer II
(A) Crystal Parameters at 20 (2) °C		
cryst dimens	0.09 × 0.12 × 0.35	0.09 × 0.51 × 0.64
cryst syst	monoclinic	monoclinic
space group	$P2_1/c$	$Cc$
<i>a</i> , Å	8.514 (2)	9.481 (3)
<i>b</i> , Å	13.577 (4)	12.477 (4)
<i>c</i> , Å	8.927 (2)	8.329 (2)
$\beta$ , deg	112.09 (2)	114.05 (2)
<i>Z</i>	4	4
<i>V</i> , Å <sup>3</sup>	956.1 (4)	900.0 (5)
<i>M<sub>r</sub></i>	128.68	128.68
$\rho_{\text{calcd}}$ , (g cm <sup>-3</sup> )	0.89	0.95
$\mu$ , cm <sup>-1</sup>	0.49	0.60
<i>F</i> (000)	288	288
(B) Data Collection		
reflens measd	$\pm h, k, l$	$h, k, \pm l$
2 $\theta$ range, deg	1–50	1–57
scan type:	$\theta$ –2 $\theta$	$\theta$ –2 $\theta$
scan speed, deg min <sup>-1</sup>	3–30	3–30
scan range, deg	1.40 + 2 $\theta$ ( $K\alpha_1$ ) to 2 $\theta$ ( $K\alpha_1$ ) – 1.25	2 $\theta$ ( $K\alpha_1$ ) + 1.3 to 2 $\theta$ ( $K\alpha_1$ ) – 1.3
total no. of reflens colld	4896	2610
no. of unique reflens colld	2199	2294
no. of obsd reflens used in refinement	702 [ $I > 4\sigma(I)$ ]	1320 [ $I > 5\sigma(I)$ ]
no. of params	82	82
<i>R<sub>F</sub></i>	0.083	0.047
<i>R<sub>wF</sub></i>	0.061	0.045
GOF	2.09	1.44

diffractometer: Syntex P3/F

radiation: Mo  $K\alpha$  ( $\lambda = 0.71069$  Å)

monochromator: highly oriented graphite cryst

bkgd measurement: stationary cryst and counter; at the beginning and end of each scan, each for one-half of the total scan time

std reflens: 3 measured every 141 reflens; no significant changes in intensities obsd [ $\bar{2}11, 0\bar{6}0, \bar{1}\bar{1}4$ ][ $\bar{3}10, 040, 043$ ]

(0.5), 128 (17), 127 (12), 126 (4), 125 (1), 115 (1), 114 (30), 113 (17), 112 (3), 111 (0.5), 110 (1), 109 (0.5), 85 (10), 84 (5), 83 (18), 82 (8), 81 (3), 80 (1), 72 (2), 71 (23), 70 (5), 69 (3), 68 (2), 67 (3), 66 (2), 57 (2), 56 (5), 55 (3), 54 (8), 53 (4), 52 (5), 51 (3), 44 (1), 43 (8), 42 (100), 41 (33), 40 (22); isomer II, *m/e* 129 [ $M^+$ ] (0.5), 128 (22), 127 (16), 126 (5), 125 (1), 124 (0.5), 123 (0.2), 115 (1), 114 (40), 113 (20), 112 (5), 111 (0.5), 110 (1), 109 (0.5), 85 (15), 84 (8), 83 (22), 82 (13), 81 (4), 80 (1), 72 (2), 71 (28), 70 (8), 69 (5), 68 (2), 67 (3), 66 (2), 57 (2), 56 (5), 55 (3), 54 (7), 53 (4), 52 (5), 51 (3), 44 (2), 43 (6), 42 (100), 41 (33), 40 (22). NMR spectra [ $(\text{CD}_3)_2\text{CO}$ ] (25 °C): <sup>11</sup>B, isomer I and II  $\delta$  – 5.4 ( $J_{\text{BH}} = 107$  Hz); <sup>1</sup>H, isomer I  $\delta$  2.98 (NH), 2.11 (BH) ( $J_{\text{BH}} = 105$  Hz), 2.09 (CH) ( $J_{\text{HH}} = 5.2$  Hz), 1.55 (BH) ( $J_{\text{BH}} = 105$  Hz); <sup>1</sup>H isomer II  $\delta$  3.43 (NH), 3.23 (NH), 3.00 (NH), 2.37 (CH) ( $J_{\text{HH}} = 6.7$ ), 2.10 (CH) ( $J_{\text{HH}} = 4.9$ ); <sup>13</sup>C, isomer I  $\delta$  34.47; <sup>13</sup>C, isomer II  $\delta$  38.33, 35.52.

**Crystallographic Measurements and Structure Solution.** The crystals were obtained as long needles, and a suitable crystal of each isomer was glued to a glass fiber. Each crystal was centered on a Syntex P3/F automated diffractometer and determinations of crystal class, orientation matrix, and unit cell dimensions were performed in a standard manner.<sup>7</sup> Data were collected in the  $\theta$ –2 $\theta$  scan mode by using Mo  $K\alpha$  radiation, a scintillation counter and a pulse-height analyzer. The data collection parameters are summarized in Table I. Examinations of the collected reflections indicated the respective space groups. The space group for isomer II suggested to be  $Cc$  from statistics was confirmed by satisfactory refinement in this space group. An empirical absorption correction based on azimuthal scans was applied for both crystals. For isomer I, the correction improved the agreement of the azimuthal scan data from 2.86% to 2.77% and the estimated maximum and minimum transmission factors were 0.879 and 0.835. For isomer II, the correction improved the agreement of the azimuthal scan data from 6.30% to 4.85% and the estimated maximum and minimum transmission factors were 0.941 and 0.721. The data were corrected for Lorentz and polarization effects,

**Table II.** Fractional Coordinates and Their Esd's for 1(e),3(e),5(e)-Trimethylcycloborazane (Isomer I) and 1(e),3(e),5(e)-Trimethylcycloborazane (Isomer II)

atom	<i>x/a</i>	<i>y/b</i>	<i>z/c</i>
Isomer I			
N(1)	0.4305 (4)	0.6960 (3)	0.0910 (4)
B(2)	0.2676 (6)	0.6365 (4)	–0.0096 (6)
C(7)	0.5836 (5)	0.6442 (4)	0.0920 (6)
N(3)	0.1078 (4)	0.6953 (3)	–0.0144 (4)
B(4)	0.0993 (7)	0.8071 (5)	0.0642 (7)
C(8)	–0.0484 (5)	0.6437 (4)	0.1161 (7)
N(5)	0.2687 (5)	0.8581 (3)	0.0403 (4)
B(6)	0.4355 (6)	0.8063 (5)	0.0443 (7)
C(9)	0.2673 (6)	0.9619 (3)	–0.0062 (6)
Isomer II			
N(1)	0.5000	0.1982 (2)	0.7500
B(2)	0.5942 (4)	0.2932 (2)	0.8675 (4)
C(7)	0.4832 (4)	0.1098 (3)	0.8592 (4)
N(3)	0.6059 (3)	0.3856 (2)	0.7436 (3)
B(4)	0.6753 (4)	0.3481 (3)	0.6068 (5)
C(8)	0.4625 (5)	0.4492 (3)	0.6605 (5)
N(5)	0.5792 (3)	0.2489 (2)	0.5012 (3)
B(6)	0.5653 (4)	0.1535 (3)	0.6181 (5)
C(9)	0.6451 (5)	0.2095 (3)	0.3781 (5)

redundant data averaged, space group extinct reflections removed, and remaining data converted to unscaled  $|F_o|$  values. The atomic scattering factors for the neutral atoms were taken from the analytical expressions given in ref 8.

All calculations were performed with the R3/SHELXLT structure determination package.<sup>9</sup> Least-squares refinements in this package use a block-cascade algorithm with full-matrix blocks of 103 parameters.<sup>10</sup> Both structures were solved by direct methods, and the best *E*-map gave trial positions for all non-hydrogen atoms. For isomer I, isotropic least-squares refinement initially on 914 reflections with  $F > 3\sigma(F)$  and 37 variables gave  $R_F = 0.24$ . Anisotropic refinement on the heavy atoms with the hydrogen atoms included in fixed idealized positions 0.96 Å from their parent atom converged at  $R_F = 0.086$  on 702 reflections with  $F > 4\sigma(F)$  and 82 variables. Anisotropic refinement in the same manner but with B–H bond distances fixed at 1.10 Å and N–H bond distances fixed at 0.85 Å (obtained from the more accurate structure determination of isomer II) led to  $R_F = 0.083$  and  $R_{wF} = 0.061$  with GOF = 2.09. The final difference map showed four peaks, which were judged to be insignificant. For isomer II isotropic least-squares refinement on 1320 reflections with  $F > 5\sigma(F)$  and 37 variables gave  $R_F = 0.134$ , and anisotropic refinement on the non-hydrogen atoms converged with  $R_F = 0.114$ . A difference map showed hydrogen atom positions for all hydrogen atoms. The hydrogen atoms were included in idealized positions 0.96 Å from their parent atoms and given fixed *U*'s equal to 1.2 times the last *U*<sub>equiv</sub> of their parent atom. The resulting agreement factors were  $R_F = 0.055$  and  $R_{wF} = 0.051$  and GOF = 1.44. The final cycles of block least-squares refinement also allowed the hydrogen atoms to vary in positions with the results  $R_F = 0.048$ ,  $R_{wF} = 0.045$ , and GOF = 1.37 for 134 parameters and 1320 reflections. The model was inverted with the same result. The final difference map showed no peak higher than 0.32 e Å<sup>-3</sup>. The final fractional coordinates are listed in Table II.


## Results and Discussion

In this study 1,3,5-trimethylcycloborazane was prepared in good yield from the combination of a commercial solution of borane-tetrahydrofuran complex and methylamine in equimolar amounts followed by reflux of a monoglyme solution of the amine–borane

(8) *International Tables for X-Ray Crystallography*; D. Reidel: Dordrecht, Holland, 1983; Vol. 1, pp 73–346.



(9) Sheldrick, G. M. *Nicolet SHELXLT Operations Manual*; Nicolet XRD: Cupertino, CA, 1981. SHELXLT uses absorption, anomalous dispersion, and scattering data compiled in: *International Tables for X-Ray Crystallography*; Kynoch: Birmingham, England, 1974; Vol. IV, pp 55–60, 99–101, 149–150.

(10) A general description of the least-squares algebra is found in: Ahmed, F. R.; Hall, S. R.; Huber, C. P., Eds. *Crystallographic Computing*; Munksgaard: Copenhagen, 1970; p 187. The least-squares refinements minimize  $\sum w(|F_o| - |F_c|)^2$  where  $w = 1/[\sigma(F_c)^2]$  for isomer I and  $w = 1/[\sigma(F_c)^2 + g(F)^2]$ ,  $g = 0.00041$  for isomer II:  $R = \sum |F_o| - |F_c| / \sum |F_o|$ ,  $R_{wF} = [\sum (|F_o| - |F_c|)^2 / \sum w|F_o|^2]^{1/2}$ , and GOF =  $[\sum w(|F_o| - |F_c|)^2 / (m - n)]^{1/2}$ , where *n* = number of parameters and *m* = number of reflections.




**Figure 1.** Infrared spectra for isomer I (top) and isomer II (bottom) recorded from KBr pellets.

complex. The course of the reaction is summarized by the eq 5 and 6. The formation of the intermediate amine-borane (98%




yield) was proven by comparison of the  $^{11}\text{B}$  NMR spectrum ( $^{11}\text{B}$ ,  $\delta -18.4$  ( $J_{\text{BH}} = 94$  Hz, quartet)) with data in the literature.<sup>11</sup> The cycloborazane was purified initially by gentle sublimation ( $70^\circ\text{C}$ ) from which a mixture of the two isomers was obtained in 69% yield. The two isomers 1(e),3(e),5(e)-trimethylcycloborazane (isomer I) and 1(e),3(e),5(a)-trimethylcycloborazane (isomer II) were separated by fractional crystallization from benzene.

Mass, infrared, and  $^1\text{H}$  and  $^{11}\text{B}$  NMR spectroscopic data have been reported, in part, in previous studies;<sup>4,5</sup> however, a more complete set of data on the separated isomers is provided in this study. Brown and Hezeltine reported that both isomers display a weak parent ion in their mass spectra, and a most intense peak at  $m/e$  114 ( $\text{M} - 15$ )<sup>+</sup>. In the present study both isomers were found to give similar spectra with a weak parent ion,  $m/e$  129, but the most intense peak appeared at  $m/e$  42 ( $\text{BNCH}_2$ )<sup>+</sup>. These results suggest that the mass spectra of **1** are very sensitive to spectrometer inlet and ion-source conditions, and care must apparently be exercised in comparing spectra of **1** obtained from different instruments. The infrared spectra (Figure 1) for the isomers are in general agreement with the spectra reported previously, and more complete listings are presented here. The  $^{11}\text{B}\{^1\text{H}\}$  NMR spectra for both isomers are identical, showing a single resonance at  $\delta -5.4$ . Restoration of the proton coupling results in broad triplets with  $J_{\text{BH}} \sim 107$  Hz. The  $^{13}\text{C}\{^1\text{H}\}$  NMR spectrum for isomer I shows a single resonance at  $\delta$  34.47, which is consistent with a single environment for the three amino methyl groups. The  $^{13}\text{C}\{^1\text{H}\}$  NMR spectrum for isomer II shows two peaks at  $\delta$  38.33 and 35.52 in approximately a 1:2 ratio. This observation is consistent with two inequivalent methyl group environments (axial and equatorial) with the axial methyl resonance appearing at lower field. The proton NMR spectra have been recorded at high field and found to agree for the most part with the assignments provided earlier.<sup>4</sup> The high-field spectra for both isomers show the N-H resonances clearly separated from the severely overlapping B-H and C-H resonances. In isomer I there is a single broad N-H resonance centered at  $\delta$  2.98 while in isomer II three resonances at  $\delta$  3.43, 3.23, and 3.00 in an approximately 1:2:0.5 ratio are observed. In the former case one N-H resonance is expected since there is only one proton environment (axial), and the broadening results from  $^{14}\text{N}$  quadrupole interactions. In the latter case a broad



**Figure 2.** Molecular geometry and atom labeling scheme for 1(e),3(e),5(e)-trimethylcycloborazane (isomer I) (25% probability ellipsoids). Selected bond distances ( $\text{\AA}$ ) and bond angles (deg): B(2)-N(1) = 1.565 (6), B(2)-N(3) = 1.564 (7), B(4)-N(3) = 1.575 (8), B(4)-N(5) = 1.556 (6), B(6)-N(1) = 1.558 (8), B(6)-N(5) = 1.573 (7), N(1)-C(7) = 1.478 (6), N(3)-C(8) = 1.476 (5), N(5)-C(9) = 1.467 (6); B(2)-N(1)-B(6) = 116.7 (3), B(2)-N(3)-B(4) = 116.5 (4), B(4)-N(5)-B(6) = 116.7 (3), N(1)-B(2)-N(3) = 109.1 (4), N(3)-B(4)-N(5) = 108.9 (4), N(1)-B(6)-N(5) = 108.7 (4), B(2)-N(1)-C(7) = 110.3 (4), B(6)-N(1)-C(7) = 110.3 (4), B(2)-N(3)-C(8) = 110.4 (4), B(4)-N(3)-C(8) = 110.4 (4), B(4)-N(5)-C(9) = 110.5 (4), B(6)-N(5)-C(9) = 110.3 (4).



**Figure 3.** Molecular geometry and atom labeling scheme for 1(e),3(e),5(a)-trimethylcycloborazane (isomer II) (25% probability ellipsoids). Selected bond distances ( $\text{\AA}$ ) and bond angles (deg): B(2)-N(1) = 1.565 (3), B(2)-N(3) = 1.582 (4), B(4)-N(3) = 1.600 (6), B(4)-N(5) = 1.576 (4), B(6)-N(1) = 1.567 (5), B(6)-N(5) = 1.577 (5), N(1)-C(7) = 1.479 (4), N(3)-C(8) = 1.480 (4), N(5)-C(9) = 1.485 (6); B(2)-N(1)-B(6) = 114.9 (2), B(2)-N(3)-B(4) = 114.3 (2), B(4)-N(5)-B(6) = 115.0 (2), N(1)-B(2)-N(3) = 108.5 (2), N(3)-B(4)-N(5) = 108.4 (3), N(1)-B(6)-N(5) = 108.7 (2), B(2)-N(1)-C(7) = 110.9 (2), B(6)-N(1)-C(7) = 109.8 (2), B(2)-N(3)-C(8) = 113.6 (3), B(4)-N(3)-C(8) = 112.9 (3), B(4)-N(5)-C(9) = 109.6 (3), B(6)-N(5)-C(9) = 109.9 (3).

two-resonance pattern (intensity ratio 1:2) is expected for the axial and equatorial N-H protons. It is assumed that the resonances at  $\delta$  3.43 and 3.23 can be assigned to these protons while the resonance at  $\delta$  3.00 is an impurity. In isomer I the inequivalent B-H protons give rise to two sets of 1:1:1:1 quartets centered at  $\delta$  2.11 ( $J_{\text{BH}} = 105$  Hz) and 1.55 ( $J_{\text{BH}} = 105$  Hz). Although B-H resonances are observed in isomer II, accurate chemical shifts for the complicated pattern have not been determined. The methyl protons in isomer I give rise to a single doublet at  $\delta$  2.09 ( $J_{\text{CH}} = 5.2$  Hz) consistent with one methyl environment while isomer II displays two doublets at  $\delta$  2.37 ( $J_{\text{HH}} = 6.7$  Hz) and 2.10 ( $J_{\text{HH}} = 4.9$  Hz) in a 1:2 area ratio. The areas indicate that the downfield doublet can be assigned to the unique axial methyl group.

Single-crystal X-ray diffraction analyses provide unambiguous confirmation of the chair-form cyclohexane-like structures for the two isomers proposed originally by Gaines and Schaeffer.<sup>4</sup> A view of the eee isomer (I) is shown in Figure 2, and a view of the eea isomer (II) is shown in Figure 3. In the eee isomer the average B-N bond length is 1.565  $\text{\AA}$ , which can be compared with the average B-N bond lengths in  $[\text{H}_2\text{NBH}_2]_3$ <sup>12</sup> (1.576  $\text{\AA}$ ),  $[\text{H}_2\text{NB-Cl}_2]_3$ <sup>13</sup> (1.575  $\text{\AA}$ ), and  $[\text{MeNBH}_2]_3$ <sup>14</sup> (1.59  $\text{\AA}$ ). The average

(11) Nöth, H.; Wrackmeyer, B. *Nuclear Magnetic Resonance Spectroscopy of Boron Compounds*; Springer-Verlag: New York, 1978. Nöth, H.; Vahrenkamp, H. *Chem. Ber.* **1966**, *99*, 1049.

(12) Corfield, P. W. R.; Shore, S. G. *J. Am. Chem. Soc.* **1973**, *95*, 1480.

terminal N-CH<sub>3</sub> distance is 1.474 Å, which compares well with the average of equatorial and axial N-CH<sub>3</sub> distances in [Me<sub>2</sub>NBH<sub>2</sub>]<sub>3</sub>, 1.49 Å. The average N-B-N angle in isomer I, 108.9°, is consistent with the anticipated tetrahedral environment for the boron atoms, and the angle is similar to the average N-B-N angles in [H<sub>2</sub>NBH<sub>2</sub>]<sub>3</sub> (107.2°) and [H<sub>2</sub>NBCl<sub>2</sub>]<sub>3</sub> (109°). The average N-B-N angle in [Me<sub>2</sub>NBH<sub>2</sub>]<sub>3</sub>, 114°, on the other hand, is considerably more open. The average B-N-B angle, 116.5°, is more open than the N-B-N angle, and it resembles the average angles in [H<sub>2</sub>NBH<sub>2</sub>]<sub>3</sub> (115.9°), [H<sub>2</sub>NBCl<sub>2</sub>]<sub>3</sub> (120.1°), and [Me<sub>2</sub>NBH<sub>2</sub>]<sub>3</sub> (113°). The B-N-C angles are essentially constant in the eee isomer at 110.4°, and this angle is related to the B-N-C angle in [Me<sub>2</sub>NBH<sub>2</sub>]<sub>3</sub>. Least-squares-planes calculations provide dihedral angles between the basal four-atom planes and the three-atom planes (three symmetry-related sets), and the B-N-B planes, 47.6°, are slightly more tilted than the N-B-N planes, 44.3°, from the B<sub>2</sub>N<sub>2</sub> planes.

The structure of the eea isomer, of course, closely resembles that of the eee isomer with the axial methyl group positioned on N(3). As expected this slightly distorts the parameters involving N(3). The average B-N bond distance is 1.578 Å with the average B-N bond distance involving N(3) being 1.591 Å and the average derived from bonds to N(1) and N(2) being 1.571 Å. On the other hand, the N-CH<sub>3</sub> distances are nearly constant, (N-C)<sub>av</sub> = 1.481 Å. The average N-B-N angle, 108.5°, compares favorably with that found in isomer I while the average B-N-B angle, 114.7°, is closed down compared to the angle in isomer I. Interestingly, the B-N-C angles for N(3) are significantly more open, (B-N-C)<sub>av</sub> = 113.4°, than the average of the remaining B-N-C angles, 110.1°. In addition, the dihedral angle between the B-(2)N(3)B(4) plane and the N(1)B(2)B(4)N(5) basal plane, 50.1°, is greater than the dihedral angle between the N(1)B(6)N(5) plane and the basal plane, 47.7°.

- (13) Hess, H.; Lux, D.; Schwarz, W. Z. *Naturforsch., B: Anorg. Chem., Org. Chem.* **1977**, *32b*, 982.  
 (14) Trefonas, L. M.; Mathews, F. C.; Lipscomb, W. N. *Acta Crystallogr.* **1961**, *14*, 273.

Although no serious attempt has been made in this study to elucidate the mechanism for the formation of 1,3,5-trimethylcycloborazane from H<sub>3</sub>B·THF and MeNH<sub>2</sub>, it is worth pointing out that two sets of experiments were performed which possibly impact on the mechanism proposed by Beachley<sup>15</sup> involving ionic four-coordinate boron species. Heating CH<sub>3</sub>NH<sub>2</sub>·BH<sub>3</sub> in monoglyme and following the reaction by <sup>11</sup>B NMR spectroscopy did not result in the observation of tricoordinate or tetracoordinate ionic boron species (δ -25 to -30 and -10 to +8) in the <sup>11</sup>B NMR spectrum. Instead, the resonance for the amine borane (δ -18.0) slowly decreased and the resonance for **1** (δ -5.4) grew in without appearance of intermediate species. Similarly, the reaction of MeNH<sub>2</sub>·HCl + LiBH<sub>4</sub> in a 1:1 ratio in monoglyme was followed by <sup>11</sup>B NMR. After 4 h at 25 °C the <sup>11</sup>B NMR spectrum showed only peaks for LiBH<sub>4</sub> (δ -40.5) and MeNH<sub>2</sub>·BH<sub>3</sub> in a 1:4 ratio. Refluxing the mixture for 2 h showed **1**, LiBH<sub>4</sub>, and MeNH<sub>2</sub>·BH<sub>3</sub> in a 4:19:1 ratio, and after 18 h only **1** and MeNH<sub>2</sub>·BH<sub>3</sub> were observed in an 18:1 ratio. Clearly, ionic species may only be present in these systems if their lifetimes are short. More detailed experiments would be required to provide further information regarding the interesting condensation process acting in the formation of **1**.

**Acknowledgment** is made to Sandia National Laboratory for support of this research under Contract 51-0623. We also acknowledge NSF Grants CHE-7802921 and CHE-8201374, which facilitated the purchases of the X-ray diffractometer and the NMR spectrometer.

**Registry No.** **1** (isomer I), 60619-50-3; **1** (isomer II), 60619-49-0; CH<sub>3</sub>NH<sub>2</sub>·BH<sub>3</sub>, 1722-33-4; H<sub>3</sub>B·THF, 14044-65-6; CH<sub>3</sub>NH<sub>2</sub>, 74-89-5.

**Supplementary Material Available:** Listings of anisotropic thermal factors and hydrogen atom parameters (4 pages); listings of observed and calculated structure factors (13 pages). Ordering information is given on any current masthead page.

- (15) Beachley, O. T. *Inorg. Chem.* **1967**, *6*, 870.

Contribution from the Departments of Chemistry, Kent State University, Kent, Ohio 44242, and Pittsburg State University, Pittsburg, Kansas 66762

## Electron Transfer. 78. Reduction of Carboxylato-Bound Chromium(V) with Bisulfite. A "Clock Reaction" Involving Chromium(IV)<sup>1</sup>

Rathindra N. Bose,\*<sup>2a</sup> N. Rajasekar,<sup>2b</sup> Dorena M. Thompson,<sup>2a</sup> and E. S. Gould\*<sup>2c</sup>

Received February 12, 1986

The chelated chromium(V) complex bis(2-ethyl-2-hydroxybutyrate)oxochromate(V) (**I**) reacts with HSO<sub>3</sub><sup>-</sup> in aqueous solutions buffered by the ligand acid, 2-ethyl-2-hydroxybutyric acid. The stoichiometry, in the presence of excess HSO<sub>3</sub><sup>-</sup>, lies close to 1:1, but the reaction passes through an intermediate, the properties of which correspond to those of a Cr(IV) complex. The latter functions as a catalyst for the primary conversion. The slow buildup of this intermediate, followed by its sudden consumption (after Cr<sup>V</sup> is depleted), results in a "clocklike" character for this reaction if monitored at or near 600 nm. The proposed mechanism for this reaction (eq 2-5 in the text) involves 1e oxidations of HSO<sub>3</sub><sup>-</sup> to the radical anion SO<sub>3</sub><sup>·-</sup> by both Cr(V) and Cr(IV) and competition for SO<sub>3</sub><sup>·-</sup> between Cr(IV) and Cr(V). Differential equations based on the proposed mechanism, in conjunction with suitably selected values for the component rate constants and for molar absorbances for the chromium species, reproduce the observed kinetic profiles. The reaction of HSO<sub>3</sub><sup>-</sup> with Cr(V) is much slower than that with the more strongly reducing Cr(IV); the rate ratio is consistent with an outer-sphere mechanism for both HSO<sub>3</sub><sup>-</sup> reactions. In contrast, the radical anion SO<sub>3</sub><sup>·-</sup> reacts with Cr(V) 20-35 times more rapidly than with Cr(IV). This inversion of relative rates, which is the basis for the autocatalysis, suggests the operation of an inner-sphere path for the Cr(V)-radical reaction.

Carboxylato chelates of chromium(V), the facile preparations of which were described in 1979,<sup>3</sup> react with an array of inorganic reductants, and these reactions exhibit considerable mechanistic

diversity. Early studies<sup>4</sup> presupposed that "even-electron" reductants would reduce Cr(V) smoothly to Cr(III), whereas 1e reductants would react through the intermediate state, Cr(IV), which, depending upon its reactivity, might or might not be detectable. It is now clear that this is an oversimplification. Reduction by hydroxylamine<sup>5</sup> yields, instead of a Cr(III) species,

- (1) Joint sponsorship of this work by the National Science Foundation (Grant No. 8313253) and by the donors of the Petroleum Research Fund, administered by the American Chemical Society, is gratefully acknowledged.  
 (2) (a) Pittsburg State University. (b) Indian Institute of Science, Bangalore, India 560012. (c) Kent State University.  
 (3) Krumpolc, M.; Roček, J. *J. Am. Chem. Soc.* **1979**, *101*, 3206.

- (4) Srinivasan, V. S.; Gould, E. S. *Inorg. Chem.* **1981**, *20*, 3176.  
 (5) Rajasekar, N.; Subramaniam, R.; Gould, E. S. *Inorg. Chem.* **1983**, *22*, 971.