

Kwantyle $f(p, n_1, n_2)$ rzędu $p = 0.95$ rozkładu Fischera-Snedecora

$n_2 \backslash n_1$	1	2	3	4	5	6	7	8	10	12	20	40	60	100	∞
1	161	200	216	225	230	234	237	239	242	244	248	251	252	253	254
2	18.5	19.0	19.2	19.2	19.3	19.3	19.4	19.4	19.4	19.4	19.4	19.5	19.5	19.5	19.5
3	10.1	9.55	9.28	9.12	9.01	8.94	8.89	8.85	8.79	8.74	8.66	8.59	8.57	8.55	8.53
4	7.71	6.94	6.59	6.39	6.26	6.16	6.09	6.04	5.96	5.91	5.80	5.72	5.69	5.66	5.63
5	6.61	5.79	5.41	5.19	5.05	4.95	4.88	4.82	4.74	4.68	4.56	4.46	4.43	4.41	4.37
6	5.99	.14	4.76	4.53	4.39	.28	.21	.15	.06	.00	3.87	3.77	3.74	3.71	3.67
7	.59	4.74	.35	.12	3.97	3.87	3.79	3.73	3.64	3.57	.44	.34	.30	.27	.23
8	.32	.46	.07	3.84	.69	.58	.50	.44	.35	.28	.15	.04	.01	2.97	2.93
9	.12	.26	3.86	.63	.48	.37	.29	.23	.14	.07	2.94	2.83	2.79	.76	.71
10	4.96	.10	.71	.48	.33	.22	.14	.07	2.98	2.91	.77	.66	.62	.59	.54
11	4.84	3.98	3.59	3.36	3.20	3.09	3.01	2.95	2.85	2.79	2.65	2.53	2.49	2.46	2.40
12	.75	.89	.49	.26	.11	.00	2.91	.85	.75	.69	.54	.43	.38	.35	.30
13	.67	.81	.41	.18	.03	2.92	.83	.77	.67	.60	.46	.34	.30	.26	.21
14	.60	.74	.34	.11	2.96	.85	.76	.70	.60	.53	.39	.27	.22	.19	.13
15	.54	.68	.29	.06	.90	.79	.71	.64	.54	.48	.33	.20	.16	.12	.07
16	.49	.63	.24	.01	.85	.74	.66	.59	.49	.42	.28	.15	.11	.07	.01
17	.45	.59	.20	2.96	.81	.70	.61	.55	.45	.38	.23	.10	.06	.02	1.96
18	.41	.55	.16	.93	.77	.66	.58	.51	.41	.34	.19	.06	.02	1.98	.92
19	.38	.52	.13	.90	.74	.63	.54	.48	.38	.31	.16	.03	1.98	.94	.88
20	.35	.49	.10	.87	.71	.60	.51	.45	.35	.28	.12	1.99	.95	.91	.84
21	4.32	3.47	3.07	2.84	2.68	2.57	2.49	2.42	2.32	2.25	2.10	1.96	1.92	1.88	1.81
22	.30	.44	.05	.82	.66	.55	.46	.40	.30	.23	.07	.94	.89	.85	.78
23	.28	.42	.03	.80	.64	.53	.44	.37	.27	.20	.05	.91	.86	.82	.76
24	.26	.40	.01	.78	.62	.51	.42	.36	.25	.18	.03	.89	.84	.80	.73
25	.24	.39	2.99	.76	.60	.49	.40	.34	.24	.16	.01	.87	.82	.78	.71
26	.23	.37	.98	.74	.59	.47	.39	.32	.22	.15	1.99	.85	.80	.76	.69
27	.21	.35	.96	.73	.57	.46	.37	.31	.20	.13	.97	.84	.79	.74	.67
28	.20	.34	.95	.71	.56	.45	.36	.29	.19	.12	.96	.82	.77	.73	.65
29	.18	.33	.93	.70	.55	.43	.35	.28	.18	.10	.94	.81	.75	.71	.64
30	.17	.32	.92	.69	.53	.42	.33	.27	.16	.09	.93	.79	.74	.70	.62
40	4.08	3.23	2.84	2.61	2.45	2.34	2.25	2.18	2.08	2.00	1.84	1.69	1.64	1.59	1.51
60	.00	.15	.76	.53	.37	.25	.17	.10	1.99	1.92	.75	.59	.53	.48	.39
120	3.92	.07	.68	.44	.29	.17	.08	.01	.91	.83	.65	.49	.42	.36	.25
∞	.84	.00	.60	.37	.21	.10	.01	1.94	.83	.75	.57	.39	.32	.24	.00

Kwantyle $f(p, n_1, n_2)$ rzędu $p = 0.975$ rozkładu Fischera-Snedecora

$n_2 \backslash n_1$	1	2	3	4	5	6	7	8	10	12	20	40	60	100	∞
1	648	800	864	900	922	937	948	957	969	977	993	1006	1010	1013	1018
2	38.5	39.0	39.2	39.2	39.3	39.3	39.4	39.4	39.4	39.4	39.4	39.5	39.5	39.5	39.5
3	17.4	16.0	15.4	15.1	14.9	14.7	14.6	14.5	14.4	14.3	14.2	14.0	14.0	14.0	13.9
4	12.2	10.6	9.98	9.60	9.36	9.20	9.07	8.98	8.84	8.75	8.56	8.41	8.36	8.32	8.26
5	10.0	8.43	7.76	7.39	7.15	6.98	6.85	6.76	6.62	6.52	6.33	6.18	6.12	6.08	6.02
6	8.81	7.26	6.60	6.23	5.99	5.82	5.70	5.60	5.46	5.37	5.17	5.01	4.92	4.92	4.85
7	.07	6.54	5.89	5.52	.29	.12	4.99	4.90	4.76	4.67	4.47	4.31	.25	.21	.14
8	7.57	.06	.42	.05	4.82	4.65	.53	.43	.30	.20	.00	3.84	3.78	3.74	3.67
9	.21	5.71	.08	4.72	.48	.32	.20	.10	3.96	3.87	3.67	.51	.45	.40	.33
10	6.94	.46	4.83	.47	.24	.07	3.95	3.85	.72	.62	.42	.26	.20	.15	.08
11	6.72	5.26	4.63	4.28	4.04	3.88	3.76	3.66	3.53	3.43	3.23	3.06	3.00	2.96	2.88
12	.55	.10	.47	.12	3.89	.73	.61	.51	.37	.28	.07	2.91	2.85	.80	.72
13	.41	4.97	.35	.00	.77	.60	.48	.39	.25	.15	2.95	.78	.72	.67	.60
14	.30	.86	.24	3.89	.66	.50	.38	.29	.15	.05	.84	.67	.61	.56	.49
15	.20	.76	.15	.80	.58	.41	.29	.20	.06	2.96	.76	.58	.52	.47	.40
16	.12	.69	.08	.73	.50	.34	.22	.12	2.99	.89	.68	.51	.45	.40	.32
17	.04	.62	.01	.66	.44	.28	.16	.06	.92	.82	.62	.44	.38	.33	.25
18	5.98	.56	3.95	.61	.38	.22	.10	.01	.87	.77	.56	.38	.32	.27	.19
19	.92	.51	.90	.56	.33	.17	.05	2.96	.82	.72	.51	.33	.27	.22	.13
20	.87	.46	.86	.51	.29	.13	.01	.91	.77	.68	.46	.29	.22	.17	.09
21	5.83	4.42	3.82	3.48	3.25	3.09	2.97	2.87	2.73	2.64	2.42	2.25	2.18	2.13	2.04
22	.79	.38	.78	.44	.22	.05	.93	.84	.70	.60	.39	.21	.14	.09	.00
23	.75	.35	.75	.41	.18	.02	.90	.81	.67	.57	.36	.18	.11	.06	1.97
24	.72	.32	.72	.38	.15	2.99	.87	.78	.64	.54	.33	.15	.08	.02	.94
25	.69	.29	.69	.35	.13	.97	.85	.75	.61	.51	.30	.12	.05	.00	.91
26	.66	.27	.67	.33	.10	.94	.82	.73	.59	.49	.28	.09	.03	1.97	.88
27	.63	.24	.65	.31	.08	.92	.80	.71	.57	.47	.25	.07	.00	.94	.85
28	.61	.22	.63	.29	.06	.90	.78	.69	.55	.45	.23	.05	1.98	.92	.83
29	.59	.20	.61	.27	.04	.88	.76	.67	.53	.43	.21	.03	.96	.90	.81
30	.57	.18	.59	.25	.03	.87	.75	.65	.51	.41	.20	.01	.94	.88	.79
40	5.42	4.05	3.46	3.13	2.90	2.74	2.62	2.53	2.39	2.29	2.07	1.88	1.80	1.74	1.64
60	.29	3.93	.34	.01	.79	.63	.51	.41	.27	.17	1.94	.74	.67	.60	.48
120	.15	.80	.22	2.89	.67	.51	.39	.30	.15	.05	.82	.61	.52	.45	.31
∞	.02	.69	.12	.79	.57	.41	.29	.19	.05	1.94	.71	.48	.39	.30	.00

Kwantyle $f(p, n_1, n_2)$ rzędu $p = 0.99$ rozkładu Fischera-Snedecora

$n_2 \backslash n_1$	1	2	3	4	5	6	7	8	10	12	20	40	60	100	∞
1	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
2	98.5	99.0	99.2	99.2	99.3	99.3	99.4	99.4	99.4	99.4	99.4	99.5	99.5	99.5	99.5
3	34.1	30.8	29.5	28.7	28.2	27.9	27.7	27.5	27.2	27.1	26.7	26.4	26.3	26.2	26.1
4	21.2	18.0	16.7	16.0	15.5	15.2	15.0	14.8	14.5	14.4	14.0	13.7	13.7	13.6	13.5
5	16.3	13.3	12.1	11.4	11.0	10.7	10.5	10.3	10.1	9.89	9.55	9.29	9.20	9.13	9.02
6	13.7	10.9	9.78	9.15	8.75	8.47	8.26	8.10	7.87	7.72	7.40	7.14	7.06	6.99	6.88
7	12.2	9.55	8.45	7.85	7.46	7.19	6.99	6.84	6.62	6.47	6.16	5.91	5.82	5.75	5.65
8	11.3	8.65	7.59	.01	6.63	6.37	.18	.03	5.81	5.67	5.36	.12	.03	4.96	4.86
9	10.6	.02	6.99	6.42	.06	5.80	5.61	5.47	.26	.11	4.81	4.57	4.48	.42	.31
10	.0	7.56	.55	5.99	5.64	.39	.20	.06	4.85	4.71	.41	.17	.08	.01	3.91
11	9.65	7.21	6.22	5.67	5.32	5.07	4.89	4.74	4.54	4.40	4.10	3.86	3.78	3.71	3.60
12	.33	6.93	5.95	.41	.06	4.82	.64	.50	.30	.16	3.86	.62	.54	.47	.36
13	.07	.70	.74	.21	4.86	.62	.44	.30	.10	3.96	.66	.43	.34	.27	.17
14	8.86	.51	.56	.04	.70	.46	.28	.14	3.94	.80	.51	.27	.18	.11	.00
15	.68	.36	.42	4.89	.56	.32	.14	.00	.80	.67	.37	.13	.05	2.98	2.87
16	.53	.23	.29	.77	.44	.20	.03	3.89	.69	.55	.26	.02	2.93	.86	.75
17	.40	.11	.18	.67	.34	.10	3.93	.79	.59	.46	.16	2.92	.83	.76	.65
18	.29	.01	.09	.58	.25	.01	.84	.71	.51	.37	.08	.84	.75	.68	.57
19	.18	5.93	.01	.50	.17	3.94	.77	.63	.43	.30	.00	.76	.67	.60	.49
20	.10	.85	4.94	.43	.10	.87	.70	.56	.37	.23	2.94	.69	.61	.54	.42
21	8.02	5.78	4.87	4.37	4.04	3.81	3.64	3.51	3.31	3.17	2.88	2.64	2.55	2.48	2.36
22	7.95	.72	.82	.31	3.99	.76	.59	.45	.26	.12	.83	.58	.50	.42	.31
23	.88	.66	.76	.26	.94	.71	.54	.41	.21	.07	.78	.54	.45	.37	.26
24	.82	.61	.72	.22	.90	.67	.50	.36	.17	.03	.74	.49	.40	.33	.21
25	.77	.57	.68	.18	.86	.63	.46	.32	.13	2.99	.70	.45	.36	.29	.17
26	.72	.53	.64	.14	.82	.59	.42	.29	.09	.96	.66	.42	.33	.25	.13
27	.68	.49	.60	.11	.78	.56	.39	.26	.06	.93	.63	.38	.29	.22	.10
28	.64	.45	.57	.07	.75	.53	.36	.23	.03	.90	.60	.35	.26	.19	.06
29	.60	.42	.54	.04	.73	.50	.33	.20	.00	.87	.57	.33	.23	.16	.03
30	.56	.39	.51	.02	.70	.47	.30	.17	2.98	.84	.55	.30	.21	.13	.01
40	7.31	5.18	4.31	3.83	3.51	3.29	3.12	2.99	2.80	2.66	2.37	2.11	2.02	1.94	1.80
60	.08	4.98	.13	.65	.34	.12	2.95	.82	.63	.50	.20	1.94	1.84	.75	.60
120	6.85	.79	3.95	.48	.18	2.96	.80	.67	.48	.34	.04	.77	.66	.56	.38
∞	.63	.61	.78	.32	.02	.80	.64	.51	.32	.18	1.88	.59	.47	.36	.00

Kwantyle $f(p, n_1, n_2)$ rzędu $p = 0.995$ rozkładu Fischera-Snedecora

$n_2 \backslash n_1$	1	2	3	4	5	6	7	8	10	12	20	40	60	100	∞
1	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
2	198	199	199	199	199	199	199	199	199	199	199	199	199	199	200
3	55.6	49.8	47.5	46.2	45.4	44.8	44.4	44.1	43.7	43.4	42.8	42.3	42.1	42.0	41.8
4	31.3	26.3	24.3	23.2	22.5	22.0	21.6	21.4	21.0	20.7	20.2	19.8	19.6	19.5	19.3
5	22.8	18.3	16.5	15.6	14.9	14.5	14.2	14.0	13.6	13.4	12.9	12.5	12.4	12.3	12.1
6	18.6	14.5	12.9	12.0	11.5	11.1	10.8	10.6	10.2	10.0	9.59	9.24	9.12	9.03	8.88
7	16.2	12.4	10.9	10.0	9.52	9.16	8.89	8.68	8.38	8.18	7.75	7.42	7.31	7.22	7.08
8	14.7	11.0	9.60	8.81	8.30	7.95	7.69	7.50	7.21	7.01	6.61	6.29	6.18	6.09	5.95
9	13.6	10.1	8.72	7.96	7.47	.13	6.88	6.69	6.42	6.23	5.83	5.52	5.41	5.32	.19
10	12.8	9.43	.08	.34	6.87	6.54	.30	.12	5.85	5.66	.27	4.97	4.86	4.77	4.64
11	12.2	8.91	7.60	6.88	6.42	6.10	5.86	5.68	5.42	5.24	4.86	4.55	4.44	4.36	4.23
12	11.8	.51	.23	.52	.07	5.76	.52	.35	.09	4.91	.53	.23	.12	.04	3.90
13	.4	.19	6.93	.23	5.79	.48	.25	.08	4.82	.64	.27	3.97	3.87	3.78	.65
14	.1	7.92	.68	.00	.56	.26	.03	4.86	.60	.43	.06	.76	.66	.57	.44
15	10.8	.70	.48	5.80	.37	.07	4.85	.67	.42	.25	3.88	.58	.48	.39	.26
16	.6	.51	.30	.64	.21	4.91	.69	.52	.27	.10	.73	.44	.33	.25	.11
17	.4	.35	.16	.50	.07	.78	.56	.39	.14	3.97	.61	.31	.21	.12	2.98
18	.2	.21	.03	.37	4.96	.66	.44	.28	.03	.86	.50	.20	.10	.01	.87
19	.1	.09	5.92	.27	.85	.56	.34	.18	3.93	.76	.40	.11	.00	2.91	.78
20	9.94	6.99	.82	.17	.76	.47	.26	.09	.85	.68	.32	.02	2.92	.83	.69
21	9.83	6.89	5.73	5.09	4.68	4.39	4.18	4.01	3.77	3.60	3.24	2.95	2.84	2.75	2.61
22	.73	.81	.65	.02	.61	.32	.11	3.94	.70	.54	.18	.88	.77	.69	.55
23	.63	.73	.58	4.95	.54	.26	.05	.88	.64	.47	.12	.82	.71	.62	.48
24	.55	.66	.52	.89	.49	.20	3.99	.83	.59	.42	.06	.77	.66	.57	.43
25	.48	.60	.46	.84	.43	.15	.94	.78	.54	.37	.01	.72	.61	.52	.38
26	.41	.54	.41	.79	.38	.10	.89	.73	.49	.33	2.97	.67	.56	.47	.33
27	.34	.49	.36	.74	.34	.06	.85	.69	.45	.28	.93	.63	.52	.43	.29
28	.28	.44	.32	.70	.30	.02	.81	.65	.41	.25	.89	.59	.48	.39	.25
29	.23	.40	.28	.66	.26	3.98	.77	.61	.38	.21	.86	.56	.45	.36	.21
30	.18	.35	.24	.62	.23	.95	.74	.58	.34	.18	.82	.52	.42	.32	.18
40	8.83	6.07	4.98	4.37	3.99	3.71	3.51	3.35	3.12	2.95	2.60	2.30	2.18	2.09	1.93
60	.49	5.80	.73	.14	.76	.49	.29	.13	2.90	.74	.39	.08	1.96	1.86	.69
120	.18	.54	.50	3.92	.55	.29	.09	2.94	.71	.55	.19	1.87	.75	.64	.43
∞	7.88	.30	.28	.72	.35	.09	2.90	.74	.52	.36	.00	.67	.53	.40	.00