

# Algebra - zadanie domowe nr 6

Zad 1 Diagonalizacja  
Baza.

$$\varphi : \mathbb{R}^3 \rightarrow \mathbb{R}^3$$

$$\varphi(x, y, z) = (x + 2y - 2z, x + 3z, x + 3y)$$

$$\det(A - \lambda I) = \begin{vmatrix} 1-\lambda & 2 & -2 \\ 1 & -\lambda & 3 \\ 1 & 3 & -\lambda \end{vmatrix} = \lambda^2(1-\lambda) - 6 + 6 - 2\lambda - 5(1-\lambda) + 2\lambda = (1-\lambda)(\lambda^2 - 9)$$

$$\text{Spec}(\varphi) = \{-3, 1, 3\} \quad \text{różne proste} \Rightarrow \varphi \text{ - diagonalizowalny}$$

•  $\lambda_1 = 1$

$$(A - \lambda_1 I)X = 0$$

$$(A - I)X = 0 \Leftrightarrow \begin{bmatrix} 0 & 2 & -2 \\ 1 & -1 & 3 \\ 1 & 3 & -1 \end{bmatrix} \begin{bmatrix} a \\ b \\ c \end{bmatrix} = \begin{bmatrix} 0 \\ 0 \\ 0 \end{bmatrix}$$

$$\begin{bmatrix} 0 & 2 & -2 \\ 1 & -1 & 3 \\ 1 & 3 & -1 \end{bmatrix} \xrightarrow{w_2 - w_1} \begin{bmatrix} 1 & -1 & 3 \\ 1 & 3 & -1 \\ 0 & 2 & -2 \end{bmatrix} \xrightarrow{w_2 - w_1} \begin{bmatrix} 1 & -1 & 3 \\ 0 & 4 & -4 \\ 0 & 2 & -2 \end{bmatrix} \xrightarrow{\substack{w_3 - w_2 \\ w_2 : 4}} \begin{bmatrix} 1 & -1 & 3 \\ 0 & 1 & -1 \\ 0 & 0 & 0 \end{bmatrix} \xrightarrow{w_1 + w_2} \begin{bmatrix} 1 & 0 & 2 \\ 0 & 1 & -1 \\ 0 & 0 & 0 \end{bmatrix}$$

$$\begin{cases} a + 2c = 0 \\ b - c = 0 \end{cases}$$

$$c = b$$

$$b = c = b$$

$$a = -2c = -2b$$

$$E_1 = \{(-2t, t, t), t \in \mathbb{R}\}$$

$$= \text{Lin}_{\mathbb{R}} \{(-2, 1, 1)\}$$

•  $\lambda_2 = 3$

$$(A - 3I)X = 0$$

$$\begin{bmatrix} -2 & 2 & -2 \\ 1 & -3 & 3 \\ 1 & 3 & -3 \end{bmatrix} \begin{bmatrix} a \\ b \\ c \end{bmatrix} = \begin{bmatrix} 0 \\ 0 \\ 0 \end{bmatrix}$$

$$\begin{bmatrix} -2 & 2 & -2 \\ 1 & -3 & 3 \\ 1 & 3 & -3 \end{bmatrix} \xrightarrow{w_1 : (-2)} \begin{bmatrix} 1 & -1 & 1 \\ 1 & -3 & 3 \\ 1 & 3 & -3 \end{bmatrix} \xrightarrow{\substack{w_2 - w_1 \\ w_3 - w_1}} \begin{bmatrix} 1 & -1 & 1 \\ 0 & -2 & 2 \\ 0 & 4 & -4 \end{bmatrix} \xrightarrow{w_2 : (-2)} \begin{bmatrix} 1 & -1 & 1 \\ 0 & 1 & -1 \\ 0 & 0 & 0 \end{bmatrix} \xrightarrow{w_1 + w_2} \begin{bmatrix} 1 & 0 & 0 \\ 0 & 1 & -1 \\ 0 & 0 & 0 \end{bmatrix}$$

$$\begin{cases} a = 0 \\ b - c = 0 \end{cases}$$

$$b = c$$

$$E_3 = \{(0, b, b), b \in \mathbb{R}\} = \text{Lin}_{\mathbb{R}} \{(0, 1, 1)\}$$

•  $\lambda_3 = -3$

$$(A + 3I)X = 0$$

$$\begin{bmatrix} 4 & 2 & -2 \\ 1 & 3 & 3 \\ 1 & 3 & 3 \end{bmatrix} \begin{bmatrix} a \\ b \\ c \end{bmatrix} = \begin{bmatrix} 0 \\ 0 \\ 0 \end{bmatrix}$$

$$\begin{bmatrix} 4 & 2 & -2 \\ 1 & 3 & 3 \\ 1 & 3 & 3 \end{bmatrix} \xrightarrow{\substack{w_3 - w_2 \\ w_1/2}} \begin{bmatrix} 2 & 1 & -1 \\ 1 & 3 & 3 \\ 0 & 0 & 0 \end{bmatrix} \xrightarrow{w_2 - w_1} \begin{bmatrix} 1 & 3 & 3 \\ 2 & 1 & -1 \\ 0 & 0 & 0 \end{bmatrix} \xrightarrow{w_2 - w_1 \cdot 2} \begin{bmatrix} 1 & 3 & 3 \\ 0 & -5 & -7 \\ 0 & 0 & 0 \end{bmatrix}$$

$$\begin{cases} a + 3b + 3c = 0 \\ -5b - 7c = 0 \end{cases}$$

$$a = -3b - 3c = \frac{21}{5}c - 3c = \frac{6}{5}c$$

$$b = -\frac{7}{5}c$$

$$c = t$$

$$E_{-3} = \{(\frac{6}{5}t, -\frac{7}{5}t, t), t \in \mathbb{R}\}$$

$$\text{Lin}_{\mathbb{R}} \left\{ \frac{1}{5}(6, -7, 5) \right\}$$

baza

$$B = \{(-2, 1, 1), (0, 1, 1), \frac{1}{5}(6, -7, 5)\}$$

$$M_{BB}(\varphi) = \begin{bmatrix} 1 & 0 & 0 \\ 0 & 3 & 0 \\ 0 & 0 & -3 \end{bmatrix}$$

Diagonalizacja endomorfizmu

Zad. 2  $\varphi: \mathbb{R}_2[x] \rightarrow \mathbb{R}_2[x] \quad \forall p \in \mathbb{R}_2[x] \quad \varphi(p)(x) = 3xp''(x) + (x^2-1)p'(1) + x \cdot p(2)$

$B_2 = (1, x, x^2)$  baza  $\mathbb{R}_2[x]$

$\varphi(1) = 3x \cdot 0 + (x^2-1) \cdot 1 + x \cdot 1 = x^2 + x - 1 = [-1, 1, 1]_{B_2}$

$\varphi(x) = 3x \cdot 0 + (x^2-1) \cdot 1 + x \cdot 2 = x^2 + 2x - 1 = [-1, 2, 1]_{B_2}$

$\varphi(x^2) = 3x \cdot 2 + (x^2-1) \cdot 1 + x \cdot 4 = x^2 + 10x - 1 = [-1, 10, 1]_{B_2}$

$A = M_{\varphi}(B_2, B_2) = \begin{bmatrix} -1 & -1 & -1 \\ 1 & 2 & 10 \\ 1 & 1 & 1 \end{bmatrix}$

$0 = \det(A - tI) = \begin{vmatrix} -1-t & -1 & -1 \\ 1 & 2-t & 10 \\ 1 & 1 & 1-t \end{vmatrix} \stackrel{W_1+W_2}{=} \begin{vmatrix} -t & 1-t & 9 \\ 1 & 2-t & 10 \\ 0 & -1+t & -9-t \end{vmatrix} \stackrel{W_1+W_3}{=} \begin{vmatrix} -t & 0 & -t \\ 1 & 2-t & 10 \\ 0 & t-1 & -9-t \end{vmatrix} =$

$= -t(2-t)(-9-t) - t(t-1) - 10(t-1)(-t) = t[(2-t)(9+t) - t+1 + 10t-10] = t[18 + 2t - 9t - t^2 + 9t - 9]$   
 $= t[-t^2 + 2t + 0]$

$\Delta = 4 + 36 = 40$   
 $\sqrt{\Delta} = 2\sqrt{10}$

$\text{Spec}(\varphi) = \{0, 1-\sqrt{10}, 1+\sqrt{10}\}$

tridno prostc

$\downarrow$   
 $\varphi$ -diagonalizowalny

Zad. 3

Niech  $\varphi \in \text{End}(\mathbb{R}^3)$  b.z.c  $\varphi(1,0,0) = (1,0,0)$   
 $\varphi(1,1,0) = (-1,-1,0)$   
 $\varphi(1,1,1) = (0,0,0)$

Oblicz  $\varphi^{100}(3,6,9)$ .

$\text{Spec}(\varphi) = \{1, -1, 0\}$

Baza wektorow wlasnych

$C = \{c_1 = (1,0,0), c_2 = (1,1,0), c_3 = (1,1,1)\}$

B-baza kanoniczna

$P = P_{B \rightarrow C} = \begin{bmatrix} 1 & 1 & 1 \\ 0 & 1 & 1 \\ 0 & 0 & 1 \end{bmatrix}$

macierz przejsci

$D = M_{CC}(\varphi) = \begin{bmatrix} 1 & 0 & 0 \\ 0 & -1 & 0 \\ 0 & 0 & 0 \end{bmatrix}$

I sposob:

$A = M_{BB}(\varphi)$

$A = P \cdot D \cdot P^{-1}, \quad A^{100} = P \cdot D^{100} \cdot P^{-1}$

$P^{-1} = ?$

$[P|I] = \left[ \begin{array}{ccc|ccc} 1 & 1 & 1 & 1 & 0 & 0 \\ 0 & 1 & 1 & 0 & 1 & 0 \\ 0 & 0 & 1 & 0 & 0 & 1 \end{array} \right] \xrightarrow{W_1-W_3} \left[ \begin{array}{ccc|ccc} 1 & 1 & 0 & 1 & 0 & -1 \\ 0 & 1 & 0 & 0 & 1 & -1 \\ 0 & 0 & 1 & 0 & 0 & 1 \end{array} \right] \xrightarrow{W_1-W_2} \left[ \begin{array}{ccc|ccc} 1 & 0 & 0 & 1 & -1 & 0 \\ 0 & 1 & 0 & 0 & 1 & -1 \\ 0 & 0 & 1 & 0 & 0 & 1 \end{array} \right] \xrightarrow{P^{-1}}$

$\varphi^{100}(3,6,9) = ?$

$A^{100} \cdot \begin{bmatrix} 3 \\ 6 \\ 9 \end{bmatrix} = P \cdot D^{100} \cdot P^{-1} \begin{bmatrix} 3 \\ 6 \\ 9 \end{bmatrix} = \begin{bmatrix} 1 & 1 & 1 \\ 0 & 1 & 1 \\ 0 & 0 & 1 \end{bmatrix} \begin{bmatrix} 1 & 0 & 0 \\ 0 & -1 & 0 \\ 0 & 0 & 0 \end{bmatrix}^{100} \begin{bmatrix} 1 & -1 & 0 \\ 0 & 1 & -1 \\ 0 & 0 & 1 \end{bmatrix} \begin{bmatrix} 3 \\ 6 \\ 9 \end{bmatrix}$

$\begin{bmatrix} 1 & 1 & 1 \\ 0 & 1 & 1 \\ 0 & 0 & 1 \end{bmatrix} \begin{bmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 0 \end{bmatrix} \cdot \begin{bmatrix} -3 \\ -3 \\ 9 \end{bmatrix} = \begin{bmatrix} 1 & 1 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 0 \end{bmatrix} \begin{bmatrix} -3 \\ -3 \\ 9 \end{bmatrix} = \begin{bmatrix} -6 \\ -3 \\ 0 \end{bmatrix}$

$\varphi^{100}(3,6,9) = (-6, -3, 0)$

II sposob

$(3,6,9) = [-3, -3, 9]_C$

$\left[ \begin{array}{ccc|c} 1 & 1 & 1 & 3 \\ 0 & 1 & 1 & 6 \\ 0 & 0 & 1 & 9 \end{array} \right] \rightarrow \left[ \begin{array}{ccc|c} 1 & 1 & 0 & -6 \\ 0 & 1 & 0 & -3 \\ 0 & 0 & 1 & 9 \end{array} \right] \rightarrow \left[ \begin{array}{ccc|c} 1 & 0 & 0 & -3 \\ 0 & 1 & 0 & -3 \\ 0 & 0 & 1 & 9 \end{array} \right]$

$D^{100} \cdot \begin{bmatrix} -3 \\ -3 \\ 9 \end{bmatrix} = \begin{bmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 0 \end{bmatrix} \begin{bmatrix} -3 \\ -3 \\ 9 \end{bmatrix} = \begin{bmatrix} -3 \\ -3 \\ 0 \end{bmatrix}$

$\varphi^{100}(3,6,9) = [-3, -3, 0]_C = -3(1,0,0) - 3(1,1,0) = (-6, -3, 0)$

Zad. 4

a)  $A = \begin{bmatrix} 1 & -5 \\ 1 & 1 \end{bmatrix} \in M_2(\mathbb{R})$

$\det(A - tI) = \begin{vmatrix} 1-t & -5 \\ 1 & 1-t \end{vmatrix} = (1-t)^2 + 5 > 0$

brak pierwiastków rzeczywistych  
 $\text{Spec}(A) = \emptyset$  A nie jest diagonalizowalna

b)  $A = \begin{bmatrix} 7 & 1 & 2 & 0 & 1 \\ 0 & 1 & 1 & 1 & 1 \\ 0 & 0 & 1 & 3 & 0 \\ 0 & 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 0 & 2 \end{bmatrix}$

A - trójkatna górna

$\text{Spec}(A) = \{ \lambda_1 = 7, \lambda_2 = 1, \lambda_3 = 2 \}$   
 $k_1 = 1, k_2 = 3, k_3 = 1$

$\dim E_{\lambda_1} = k_1 = 1$

$\dim E_{\lambda_3} = k_3 = 1$

Zai  $\dim E_{\lambda_2} = 5 - r(A - 1 \cdot I) = 5 - 4 = 1 \neq k_2 = 3$

$A - I = \begin{bmatrix} 6 & 1 & 2 & 0 & 1 \\ 0 & 0 & 1 & 1 & 1 \\ 0 & 0 & 0 & 3 & 0 \\ 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 1 \end{bmatrix}$

$r(A - I) = 4$

NIE jest diagonalizowalna

c)  $A = \begin{bmatrix} 2 & 2 & 2 & 2 \\ 0 & 0 & 0 & 0 \\ 3 & 3 & 3 & 3 \\ 2 & 2 & 2 & 2 \end{bmatrix}$

$\lambda_1 = 0 \in \text{Spec}(A)$   
 $k_1 = 3$

$\dim E_{\lambda_1} = ?$

$\dim E_{\lambda_1} = 4 - r(A - 0 \cdot I) = 4 - r(A) = 4 - 1 = 3 = k_1$

$r(A) = 1$  rząd!

TAK, diagonalizowalna

d)  $A = \begin{bmatrix} 5 & -2 & 6 & -1 \\ 0 & 3 & p & 0 \\ 0 & 0 & 5 & 4 \\ 0 & 0 & 0 & 1 \end{bmatrix}$

dla jakich p macierz jest diagonalizowalna?

$\text{Spec}(A) = \{ \lambda_1 = 5, \lambda_2 = 3, \lambda_3 = 1 \}$   
 $k_1 = 2, k_2 = 1, k_3 = 1$

Wz. mozy tw. spektralnego, A diagonalizowalna  $\Leftrightarrow \dim E_{\lambda_1} = 2$

$\dim E_{\lambda_1} = 4 - r(A - 5I) = 2 \Leftrightarrow r(A - 5I) = 4 - 2 = 2$

$r(A - 5I) = r \begin{bmatrix} 0 & -2 & 6 & -1 \\ 0 & -2 & p & 0 \\ 0 & 0 & 0 & 4 \\ 0 & 0 & 0 & -4 \end{bmatrix} = r \begin{bmatrix} 0 & -2 & 6 & -1 \\ 0 & -2 & p & 0 \\ 0 & 0 & 0 & 1 \end{bmatrix} = r \begin{bmatrix} -2 & 6 & -1 \\ -2 & p & 0 \\ 0 & 0 & 1 \end{bmatrix} \stackrel{w_2 - w_1}{=} r \begin{bmatrix} -2 & 6 & -1 \\ 0 & p-6 & 1 \\ 0 & 0 & 1 \end{bmatrix}$

$r(A - 5I) = \begin{cases} 3 & ; p \neq 6 \\ 2 & ; p = 6 \end{cases}$

A diagonalizowalna  $\Leftrightarrow p = 6$

Zad. 5 PRAWDA / FAŁSZ

$A = \begin{bmatrix} 7 & -3 & -1 & 0 \\ 0 & 4 & p^3 & 0 \\ 0 & 0 & 7 & \pi \\ 0 & 0 & 0 & 1 \end{bmatrix}$

$A = M_{\mathcal{U}}(B_{\mathcal{U}}^4, B_{\mathcal{U}}^4)$

$A - 7I = \begin{bmatrix} 0 & -3 & -1 & 0 \\ 0 & -3 & p^3 & 0 \\ 0 & 0 & 0 & \pi \\ 0 & 0 & 0 & 1 \end{bmatrix}$

$k_1 = 2, k_2 = 1, k_3 = 1$   
 $\bullet \text{Spec}(A) = \{ \lambda_1 = 7, \lambda_2 = 4, \lambda_3 = 1 \} \Rightarrow A \text{ diagonalizowalna} \Leftrightarrow \dim E_{\lambda_1} = 4 - r(A - 7I) = 2$   
 $\Leftrightarrow r(A - 7I) = 2$

$\bullet \det A = 7 \cdot 4 \cdot 7 = 196 \neq 0 \Rightarrow r(A) = r(\mathcal{U}) = 4 \Rightarrow \mathcal{U} \text{ surjeczna}$   
 $\Rightarrow \dim \text{Ker } \mathcal{U} = 4 - 4 = 0 \Rightarrow \mathcal{U} \text{ iniekcja}$

$\bullet \det(\frac{1}{7} A^T) = \frac{1}{7^2} \cdot 196$

$r(A - 7I) = r \begin{bmatrix} 0 & -3 & -1 & 0 \\ 0 & 0 & p^3 + 1 & 0 \\ 0 & 0 & 0 & 1 \end{bmatrix} = \begin{cases} 2 & ; p = -1 \\ 3 & ; p \neq -1 \end{cases}$

$\downarrow$   
 a) ~~FAŁSZ~~ PRAWDA  
 b) FAŁSZ  
 c) PRAWDA