

L. PASICKI (Lublin)

### On the measures of non-compactness

In [1] K. Kuratowski defined a measure of non-compactness and proved the theorem of non-empty intersection for it.

We give here a shorter proof of this theorem.

**THEOREM 1.** *Let  $(M, d)$  be a complete metric space and let  $b: 2^M \rightarrow \bar{R}$  be a measure of non-compactness satisfying the following conditions:*

- 1°  $b(E) = 0$  iff  $\bar{E}$  is compact,
- 2° if  $E \subset F$ , then  $b(E) \leq b(F)$ ,
- 3°  $b(E) = b(E \cup \{x\})$  for  $x \in M$ .

*Then, if for a chain  $(\mathcal{F}, C)$  of closed non-empty subsets of  $M$  we have  $\inf\{b(F) : F \in \mathcal{F}\} = 0$ , then  $\bigcap_{F \in \mathcal{F}} F \neq \emptyset$ .*

**Proof.** Let us consider a subsequence  $(F_n)_{n \in N}$  such that, for all  $n \in N$ ,  $F_{n+1} \subset F_n$  and  $\lim_{n \rightarrow \infty} b(F_n) = 0$ . We can choose elements  $x_n \in F_n$ . Let us consider the set  $D = \bigcup_{n=1}^{\infty} \{x_n\}$ . By 3° for all  $k \in N$  we have

$$b(D) = b\left(\bigcup_{n=1}^{\infty} \{x_n\}\right) = b\left(\bigcup_{n=1}^{\infty} \{x_n\}\right) \leq b(F_k)$$

and therefore  $b(D) = 0$ . All the limits of the convergent subsequences of  $(x_n)_{n \in N}$  are contained in  $F := \bigcap_{n=1}^{\infty} F_n$ , which is closed and compact. Now we may consider only a chain consisting of compact sets, which, as it is known, have a non-empty intersection. Q.E.D.

Let  $(X, T)$  be a linear topological space.

**DEFINITION 1.** We say that for  $Y \subset X$  a mapping  $f: Y \rightarrow Y$  is *generalized condensing* if for  $A \subset Y$  and such that  $f(A) \subset A$  from the compactness of  $A - \overline{\text{co}} f(A)$  it follows that  $\bar{A}$  is compact (cf. [2]).

We present below a modification of the result obtained in [2].

**THEOREM 2.** *If  $A = \overline{\text{co}} A \subset X$  and  $f: A \rightarrow A$  is a generalized condensing mapping, then there exists a compact convex set  $C$  such that  $\overline{\text{co}} f(C) = C$ .*

**LEMMA.** *If  $A = \overline{\text{co}} A$ ,  $f: A \rightarrow A$  and there exists a compact set  $B \subset A$  such that  $f(B) \subset B$ , then there exists a  $C = \overline{\text{co}} C \subset A$  such that  $\overline{\text{co}} f(C) = C$ .*

**Proof.** Let us consider a family  $\mathcal{F} = \{F = \overline{\text{co}} F \subset A: \overline{\text{co}} f(F) \subset F\}$ . It is non-empty, as  $A \in \mathcal{F}$ . By the Hausdorff theorem there exists a maximal chain  $\mathcal{H} \subset \mathcal{F}$  containing a set that has a non-empty intersection with  $B$ . Let  $\mathcal{G} \subset \mathcal{H}$  consist of all sets  $G$  such that  $G \cap B \neq \emptyset$ . The set  $C := \bigcap_{G \in \mathcal{G}} G$  is non-empty because, by the compactness of  $B$ ,  $\bigcap_{G \in \mathcal{G}} G \cap B \neq \emptyset$ . Moreover,  $C = \overline{\text{co}} C$ ,  $\overline{\text{co}} f(C) \subset C$  and  $\overline{\text{co}} f(C) \in \mathcal{G}$ , as  $f(B \cap C) \subset B$ . By virtue of the definition of  $C$  it must be  $\overline{\text{co}} f(C) = C$ .

**Proof of the theorem.** In view of the lemma and the definition of the generalized condensing mapping, the theorem will be proved if we find a compact set  $B$  such that  $f(B) \subset B$ .

Let  $D$  be any compact subset of  $A$ , and  $B = \overline{B}$  the minimal set containing  $D$  such that  $f(B) \subset B$ . We see that  $B - \overline{f(B)} \subset D$  because  $B - \overline{f(B)} \cap (X - D)$  is relatively open in  $B$  and would be rejected when non-empty. Now, by the definition of  $f$ ,  $B$  is compact.

**COROLLARY.** *If  $(X, T)$  is a topological space and for a mapping  $f: X \rightarrow X$  there exists a compact set  $B$  such that  $f(B) \subset B$ , then we can find a compact set  $C \subset X$  for which  $f(C) = C$ .*

**Proof.** We may follow the procedure that was used in the proof of our lemma, with substituting the operation “ $\overline{\text{co}}$ ” by “ $\overline{\quad}$ ” (closure), and considering a maximal chain containing  $B$ .

**DEFINITION 2.** We say that a mapping  $f: X \rightarrow X$  is *b-condensing* ( $b: 2^X \rightarrow \overline{\mathbb{R}}$ ) if for all  $A \subset X$  such that  $b(f(A)) < \infty$  and  $b(A) \leq b(f(A))$ ,  $\overline{f(A)}$  is compact.

We see that if  $b$  satisfies  $1^\circ$ ,  $2^\circ$  and

$$3^\circ \quad b(E) = b(\overline{\text{co}} E \cup Z) \text{ for all } Z \subset A \text{ and } b(Z) = 0,$$

then for  $Y \subset X$  and  $b(Y) < \infty$ , any  $b$ -condensing mapping  $f: Y \rightarrow Y$  is generalized condensing.

Now, let  $(B, \| \cdot \|)$  be a Banach space and let  $s: 2^B \rightarrow \overline{\mathbb{R}}$  denote the Hausdorff measure of non-compactness (for  $A \subset B: s(A) = \inf \{r \in \overline{\mathbb{R}}: \text{there exists a finite } r\text{-net in } A\}$ ). It is known that for  $E, F \subset B$   $s$  satisfies  $1^\circ$  and the following conditions:

$$4^\circ \quad s(E \cup F) = \max \{s(E), s(F)\},$$

$$5^\circ \quad s(\overline{\text{co}} E) = s(E),$$

$6^\circ$  if  $N_e(E) := \{x \in B: \text{dist}(x, E) \leq e\}$ , then there exists a function  $k: \mathbb{R}^+ \rightarrow \overline{\mathbb{R}}$  such that  $\lim_{e \rightarrow 0^+} k(e) = 0$ ,  $s(N_e(E)) \leq s(E) + k(e)$ .

THEOREM 3. The measure of non-compactness  $b: 2^B \rightarrow \bar{R}$  satisfies  $1^\circ, 4^\circ-6^\circ$  iff there exists a non-decreasing continuous function  $g: \bar{R} \rightarrow \bar{R}$  such that  $b = g \circ s$ .

Proof. (i) For  $x \in B, r \in R^+$  and the balls  $K \subset B$  we have  $\lim_{e \rightarrow 0} b(K(x, r+e)) = b(K(x, r))$ .

Let  $0 < e$ ; then in view of  $4^\circ$  and, on the other hand, by  $6^\circ$ ,

$$b(K(x, r)) \leq b(K(x, r+e)) \leq b(K(x, r)) + k(e),$$

and similarly, for  $-r \leq e < 0$ ,

$$b(K(r-e)) \leq b(K(x, r)) \leq b(K(x, r-e)) + k(e)$$

which proves (i).

(ii) For all  $x \in B, r \in R^+ b(K(x, r)) = b(K(0, r))$ .

For each  $0 < e < r$  we can choose  $t \in R^+$  such that  $K(x, r-e) \subset \bar{co} \{K(0, r) \cup \{tx\}\}$  and therefore by  $4^\circ$  and  $5^\circ, b(K(x, r-e)) \leq b(K(0, r))$  and similarly  $b(K(0, r-e)) \leq b(K(x, r))$ . Now (ii) is a consequence of (i).

(iii) Let  $E \subset B, s(E) = r$ . From the definition of  $s$  and by  $4^\circ, (ii)$  and  $6^\circ$  we have:

$$b(E) = b\left(\bigcup_{i \in I_e} K(x_i, r+e)\right) = b(K(0, r+e)) \leq b(E) + k(e),$$

where  $E \subset \left(\bigcup_{i \in I_e} K(x_i, r+e)\right)$  and  $I_e$  is finite. Therefore

$$b(E) = b(K(0, r)) =: g(r) = g \circ s(K(0, r)) = g \circ s(E)$$

which with (i) and  $4^\circ$  gives the thesis of our theorem.

COROLLARY. If  $f: B \rightarrow B$  is a  $b$ -condensing mapping and  $b$  satisfies  $1^\circ, 4^\circ-6^\circ$ , then  $f$  is  $s$ -condensing.

Proof. Obviously, if  $s(f(A)) < s(A)$ , then  $b(f(A)) \leq b(A)$  and equivalently, because of the properties of  $g$ , from  $b(A) \leq b(f(A))$  it follows that  $s(A) \leq s(f(A))$ .

#### References

- [1] K. Kuratowski, *Sur les espaces complets*, Fund. Math. 15 (1930), p. 301-309.  
 [2] E. A. Lifschic, B. N. Sadovskij, Theory of fixed point for generalized condensing operators (in Russian). D.A.N. SSSR 183.2 (1968), p. 278-279.