

## RETRACTS IN METRIC SPACES

LECH PASICKI<sup>1</sup>

**ABSTRACT.** In this paper we define  $S$ -contractibility and two classes of spaces connected with this notion. A space  $X$  is said to be  $S$ -contractible provided that  $S$  is a function  $S: X \times \langle 0, 1 \rangle \times X \ni (x, \alpha, y) \mapsto S_x(\alpha, y) \in X$  that is continuous in  $\alpha$  and  $y$ , and for every  $x, y \in X$ ,  $S_x(0, y) = y$ ,  $S_x(1, y) = x$ . This notion is close to equiconnectedness, which can be defined as follows. A space  $X$  is equiconnected if there exists a map  $S$  such that  $X$  is  $S$ -contractible and  $S_x(\alpha, x) = x$  for all  $x \in X$  and  $\alpha \in I$  (cf. [4]). The results we obtain in the theory of retracts are close to those that are known for equiconnected spaces. Also the thickness of the neighborhood that can be retracted on a set in a metric space is estimated, which enables to prove a theorem belonging to fixed point theory.

1. We repeat the notions related to equiconnectedness [2].

**DEFINITIONS.** A *local equiconnecting function* for a space  $X$  is a map  $\lambda: U \times I \rightarrow X$ , where  $U$  is a neighborhood of the diagonal in  $X \times X$  such that  $\lambda(x_0, x_1, i) = x_i$ ,  $i = 0, 1$ , and  $\lambda(x, x, t) = x$  for every  $x_0, x_1, x \in X$ ,  $t \in I$ .

The  $\lambda$ -*extension* of a subset  $A \subset X$  is the smallest nonempty subset  $\hat{A} \subset X$  (if it exists) such that  $A \times \hat{A} \subset U$  and  $\lambda(A \times \hat{A} \times I) \subset \hat{A}$ .  $A$  is  $\lambda$ -*convex* if  $A = \hat{A}$ .

A local equiconnecting function  $\lambda$  is *stable* if for every neighborhood  $N$  of any point  $p \in X$  there exists a neighborhood  $M$  such that  $\hat{M} \subset N$  [3].

For  $\mathcal{U}$  an open cover of  $X$  and  $n \geq 1$  let  $X^n(\mathcal{U}) = \{(x_1, \dots, x_n) \in X^n: \{x_1, \dots, x_n\} \subset U \in \mathcal{U}\}$  with the relative topology. Let  $T^{n-1}$  denote the standard  $(n-1)$  simplex in Euclidean  $n$ -space:  $T^{n-1} = \{(t_1, \dots, t_n) \in R^n: t_i > 0, \sum t_i = 1\}$ .

A *local convex structure* for a space  $X$  consists of an open cover  $\mathcal{U}$  and a sequence of maps  $\lambda^n: X^n(\mathcal{U}) \times T^{n-1} \rightarrow X$ ,  $n \geq 1$ , such that

(i)  $\lambda^n(x_1, \dots, x_n; t_1, \dots, t_n) = \lambda^{n-1}(x_1, \dots, \bar{x}_m, \dots, x_n; t_1, \dots, \bar{t}_m, \dots, t_n)$  if  $t_m = 0$ ,

(ii) for every neighborhood  $N$  of any point  $p \in X$  there exists a neighborhood  $M$  such that  $\lambda^n(M^n \times T^{n-1}) \subset N$  for all  $n$  [5].

$X$  is called *stably LEC* if it admits a local equiconnecting function, and  $X$  is *LCS* if it admits a local convex structure.

If such a map  $\lambda$  is defined on the whole  $X \times X \times I$  then  $X$  is *stably EC* or *CS* respectively.

Received by the editors April 23, 1978 and, in revised form, March 21, 1979.

AMS (MOS) subject classifications (1970). Primary 54C15.

Key words and phrases. Retraction, metric space, contractible set.

<sup>1</sup>The author is indebted to the referee for his valuable remarks.

© 1980 American Mathematical Society  
 0002-9939/80/0000-0178/\$02.50

2. DEFINITION 1. Let a set and a function  $S$  be given that satisfy the following conditions:

$$(1) S: X \times I \times X \ni (x, t, y) \mapsto S_x(t, y) \in X,$$

$$(2) S_x(0, y) = y, S_x(1, y) = x \text{ for any } x, y \in X.$$

Then for any nonempty set  $A \subset X$  let  $\text{coS } A = \inf\{D \subset X: A \subset D \text{ and for any } x \in A, t \in I, S_x(t, D) \subset D\}$ . For  $A = \emptyset$  let  $\text{coS } A = \emptyset$ . If  $\text{coS } A = A$  then  $A$  is  $S$ -convex.

The above definition is correct (i.e. the infimum exists) because for any two sets  $E, D$  such that, for any  $x \in A$  and  $t \in I, S_x(t, D) \subset D$  and  $S_x(t, E) \subset E$  we have  $S_x(t, D \cap E) \subset D$  and  $S_x(t, D \cap E) \subset E$  which implies  $S_x(t, D \cap E) \subset D \cap E$ .

PROPOSITION 1. If  $\{A_s\}_{s \in T}$  is a family of  $S$ -convex sets, then  $\bigcap_{s \in T} A_s$  is  $S$ -convex.

PROOF. Suppose that  $\bigcap_{s \in T} A_s \neq \emptyset$ . For any  $x \in \bigcap_{s \in T} A_s, t \in I$  and  $s \in T$  we have that  $S_x(t, \bigcap_{s \in T} A_s) \subset A_s$  and consequently  $S_x(t, \bigcap_{s \in T} A_s) \subset \bigcap_{s \in T} A_s$ , which means that  $\bigcap_{s \in T} A_s$  is  $S$ -convex.

DEFINITION 2. A space is  $S$ -contractible if  $S$  satisfies the conditions (1), (2), and, for any  $x \in X, \{S_x(t, \cdot)\}$  is a homotopy joining the identity with a constant map (cf. [1, p. 22]).

DEFINITION 3. A space  $X$  is of  $C$  type I if  $C$  is a subset of  $X$  and there exists  $S$  such that  $X$  is  $S$ -contractible and

(3) for any  $x \in C$  and any neighborhood  $N$  of  $x$  there exists a neighborhood  $U$  such that  $\text{coS } U \subset N$ .

If  $C = X$  then we say it is of type I.

Obviously any stably EC is of type I.

Let  $(M, d)$  be a metric space. For the nonempty sets  $A, D \subset M$  and  $r > 0$  let us write  $d(A, D) = \inf\{d(x, y): x \in A, y \in D\}, B(A, r) = \{x \in M: d(A, x) < r\}$  and  $\text{dia } A = \sup\{d(x, y): x, y \in A\}$ .

THEOREM 1. Let  $(M, d)$  be a metric space and let  $A = \bar{A}$  be of  $\partial A$  type I ( $\partial A$  denotes the boundary of  $A$ ) such that, for any  $x \notin A, d(x, A) = d(x, \partial A)$ . Then  $A$  is retract of  $M$ .

PROOF. Let  $\{U_s\}_{s \in T}$  be a locally finite open cover of  $M \setminus A$  with a well-ordered family of indices  $T$  and, for  $\{a_s\}_{s \in T} \subset \partial A$ , let the following condition be satisfied: if  $x \in U_s$ , then  $d(x, a_s) \leq 2d(x, A)$  for  $s \in T$  ([1, p. 70]).

For  $x \in M \setminus A$  let us consider  $T_x := \{s \in T: x \in U_s\}$  and let

$$c_s(x) = d(x, M \setminus U_s) / \sup\{d(x, M \setminus U_s): s \in T_x\} \quad (4)$$

and let

$$r(x) = \begin{cases} x & \text{for } x \in A, \\ S_{a_1}(c_{s_1}(x), S_{a_2}(c_{s_2}(x), \dots, (S_{a_n}(c_{s_n}(x), y) \dots))) & \text{for } x \in M \setminus A, \end{cases} \quad (5)$$

where  $\{s_1, s_2, \dots, s_n\} = T_x$  and  $s_1 < s_2 < \dots < s_n$  and  $y \in A$ .

It is easily seen that there always exists  $s \in T_x$  such that  $c_s(x) = 1$ ; then  $S_{a_s}(c_s(x), z) = a_s$  for  $z \in A$ . It is trivial that  $r(x) \in \text{coS}\{B(x, 2d(x, A)) \cap A\}$ . So it follows from (3) that  $r$  is continuous on  $\partial A$ . Also  $r$  is continuous on  $M \setminus A$  as for any  $x \in M \setminus A$  there exists  $B(x, \delta(x))$  which meets only finitely many  $U_s$ . Then  $T_z$  is finite and fixed for  $z \in B(x, \delta(x))$ , and  $r$  is a finite superposition of the same continuous maps in  $B(x, \delta(x))$ .

**PROPOSITION 2.** *Any metric space which is of type I is an AR( $\mathcal{U}$ ).*

**COROLLARY 1.** *Any metrizable space which is of type I is CS (cf. [5]).*

**DEFINITION [1, p. 219].** A compact space  $X$  that is metrizable in such a way that for any  $x, y \in X$  there exists exactly one  $z$  such that  $\rho(x, z) = \rho(y, z) = \rho(x, y)/2$  is called a *strongly convex compactum*.

**COROLLARY 2.** *Any strongly convex compactum is AR (cf. [1, p. 219]).*

**DEFINITION 4.** A space  $X$  is of  $C$  type II provided that  $C \subset X$  and there exists  $S$  such that  $X$  is  $S$ -contractible and the following condition holds:

(6) for any neighborhood  $N$  of any  $x \in C$  there exists a neighborhood  $U$  such that for every  $z \in U \cap C$  and  $t \in I$  we have  $S_z(t, U) \subset N$ .

If  $C = X$  let us call it type II.

It is easily seen that every type I is type II.

If  $X$  is a locally compact space which is  $S$ -contractible and  $S$  is a map, and if  $S_x(t, x) = x$  for all  $x \in X$ , then  $X$  is of type II.

**PROPOSITION 3.** *Let  $A = \bar{A}$  be a  $\partial A$  type II subset of a metric space  $(M, d)$  such that, for any  $x \notin A$ ,  $d(x, A) = d(x, \partial A)$  and  $M \setminus A$  is finite dimensional. Then  $A$  is retract of  $M$ .*

**PROOF.** Let  $\dim M \setminus A \leq n$ . Then we may assume that every  $x \in M \setminus A$  belongs to at most  $n + 1$  sets of  $\{U_s\}_{s \in T}$  and we follow the proof of Theorem 1. Condition (6) then ensures the continuity of  $r$  on  $\partial A$ .

**THEOREM 2.** *Let  $A = \bar{A}$  be a  $\partial A$  type II subset of a finite dimensional subspace of a linear normed space  $(X, \|\cdot\|)$ . Then  $A$  is a retract of  $X$ .*

**PROOF.** We construct a dense set  $E$  in  $\partial A$  in a special way.

1°. Let  $E_1 \subset \partial A \cap B(0, 1)$  be a minimal set with respect to the property that for every  $x \in \partial A \cap B(0, 1)$ ,  $d(x, E_1) \leq 1$ . We denote the elements of  $E_1$  by the natural numbers.

2°. We complete  $E_1$  to  $E_2 \subset \partial A \cap B(0, 2)$  a minimal set with respect to the property that for any  $x \in \partial A \cap B(0, 2)$ ,  $d(x, E_2) \leq 1/2$  and sign "new" points by the further numbers, etc.

$n^\circ$ . We complete  $E_{n-1}$  to  $E_n \subset \partial A \cap B(0, n)$ ; for  $x \in \partial A \cap B(0, n)$ ,  $d(x, E_n) \leq 1/n$ .

Now let  $E = \bigcup_{n=1}^\infty E_n$  and for  $a_n \in E$  let

$$c_n(x) = \max\{0, \min\{1, 3 - d(x, a_n)/d(x, A)\}\} \tag{7}$$

and for  $x \in X, y \in A$ ,

$$\begin{aligned}
 p_1(x, y) &= S_{a_1}(c_1(x), y), \\
 p_n(x, y) &= p_{n-1}(x, S_{a_n}(c_n(x), y)) \quad \text{for } n > 1.
 \end{aligned}
 \tag{8}$$

We define  $r: X \rightarrow A$  as follows:

$$r(x) = \begin{cases} x & \text{for } x \in A, \\ \lim_{n \rightarrow \infty} p_n(x, y) & \text{for } x \notin A. \end{cases}
 \tag{9}$$

The set  $A$  is contained in a finite dimensional subspace of  $X$ , which with the linearity of norm yields that for  $x \in B(0, r)$  each of the sets  $A \cap B(x, 2d(x, A))$  and  $A \cap B(x, 3d(x, A)) \setminus B(x, 2d(x, A))$  contains at least one and not more than  $k$  elements of the sets  $E_{n(x)}$ , where  $n(x) = \max\{\lceil 8/d(x, A) \rceil, \lceil (r + 1)/6 \rceil\}$ . In view of the construction of  $E$  we need not consider the superposition of more than  $k$  maps because there are at most  $m \leq k$  coefficients  $c_n(x) \in (0, 1)$  before the first one that is equal to 1. Therefore  $r$  is continuous on  $X \setminus A$ . The continuity on  $\partial A$  follows from (6).

DEFINITION 5. A space  $X$  is locally  $S$ -contractible if there exists  $S$  satisfying (1), (2) and

(10) for any  $x \in X$  there exists a neighborhood  $U$  such that, for any  $z \in U$ ,  $\{S_z(t, \cdot)\}|_U$  is a homotopy joining the identity with a constant map.

DEFINITION 6. A space  $X$  is locally  $C$  type I ( $C$  type II) if it is locally  $S$ -contractible and (3) ((6)) is satisfied.

It is obvious that every LEC space is locally  $S$ -contractible and every stably LEC space is locally type I.

THEOREM 3. Let  $A = \bar{A}$  be locally  $A$  type I in a metric space  $(M, d)$  such that, for every  $x \notin A$ ,  $d(x, A) = d(x, \partial A)$ . Then  $A$  is a retract of  $D$ , if  $D$  is as follows.

$$\begin{aligned}
 D = \{x \in M: \text{there exists } \epsilon > 0 \text{ such that, for } y, z \in \text{coS}\{B(x, d(x, A) + \epsilon) \cap \partial A\} \\
 \text{and } z \in \partial A, S_z \text{ is a map}\}.
 \end{aligned}
 \tag{11}$$

PROOF. Let  $\epsilon(x) = \sup\{\epsilon: \text{such that for } y, z \in \text{coS}\{B(x, d(x, A) + \epsilon) \cap \partial A\} \text{ and } z \in \partial A, S_z \text{ is a map}\}$  and let  $(x_n)_{n \in N}$  be any sequence convergent in  $D$ , say to  $x_0$ . We have  $\lim_{n \rightarrow \infty} \epsilon(x_n) \leq \epsilon(x_0)$  because otherwise there would exist  $n \in N$  and  $\delta > 0$  such that  $B(x_0, d(x_0, A) + \epsilon(x_0) + \delta) \subset B(x_n, d(x_n, A) + \epsilon(x_n))$ . Similarly  $\epsilon(x_0) \leq \liminf_{n \rightarrow \infty} \epsilon(x_n)$  because otherwise  $B(x_n, d(x_n, A) + \epsilon(x_n) + \delta) \subset B(x_0, d(x_0, A) + \epsilon(x_0))$  would hold for a  $\delta > 0$ . Now let  $D_\delta = \text{Int}\{x \in D: \epsilon(x) \geq \delta\}$  and  $\mathfrak{B} = \{B(x, \lambda(x)) \cap D_{\lambda(x)}: x \in D \setminus A\}$  where

$$\lambda(x) = \min\{d(x, A), \epsilon(x)/4\}.
 \tag{12}$$

If for  $x \in D \setminus A$  there exists  $\delta > 0$  such that, for each  $y \in B(x, \delta)$ ,  $\lambda(y) \geq \lambda(x)$  then  $x \in D_{\lambda(x)}$ , otherwise a  $y$  can be found such that  $\lambda(y) < \lambda(x)$  (implies  $x \in D_{\lambda(y)}$ ) and  $x \in B(y, \lambda(y))$ . Hence  $\mathfrak{B}$  is an open cover of  $D \setminus A$  and we can find a

locally finite open cover  $\{U_s\}_{s \in T}$  which is a star refinement of  $\mathfrak{B}$ . If for  $s \in T$ ,  $x_s \in U_s$ , we choose  $z$  for which  $\text{St}(U_s, \mathcal{U}) \subset B(z, \lambda(z)) \cap D_{\lambda(z)}$  and  $a_s \in B(x_s, d(x_s, A) + \lambda(z)) \cap \partial A$ , then for  $x \in U_s$  we have

$$\begin{aligned} d(x, a_s) &\leq d(x, x_s) + d(x_s, a_s) \leq \lambda(z) + d(x_s, A) + \lambda(z) \\ &\leq 2\lambda(z) + d(x_s, x) + d(x, A) \leq 3\lambda(z) + d(x, A) \\ &< d(x, A) + \varepsilon(x). \end{aligned}$$

Now it is easily seen that for these  $\{U_s\}_{s \in T}$  and  $\{a_s\}_{s \in T}$  formulas (4) and (5) give the required retraction of  $D$ .

**PROPOSITION 4.** *Any metric space which is locally type I is an ANR ( $\mathcal{U}$ ).*

**PROOF.** In the previous considerations we put everywhere "A" in place of " $\partial A$ ". We see that  $D' = \bigcup_{\delta > 0} D'_\delta$  so obtained is open. If  $x \in A$  then there is  $\delta > 0$  for which  $x \in D'$  and hence  $A \subset D'$ .

**COROLLARY.** *Any metrizable locally type I space is LCS (cf. [5]).*

**PROPOSITION 5.** *Let  $A = \bar{A}$  be locally A type I in a metric space  $(M, d)$  such that, for every  $x \notin A$ ,  $d(x, A) = d(x, \partial A)$  and  $\inf\{\sup\{r: S_z \text{ is a map for } z, y \in \text{coS}\{B(x, r) \cap \partial A\} \text{ and } z \in \partial A\}: x \in \partial A\} = a > 0$ . Then  $A$  is a retract of  $B(A, a/2)$ .*

**PROOF.** It is enough to show that  $B(A, a/2) \subset D$ , where  $D$  is defined by (11). Let  $x \in B(A, a/2)$ . Then  $\delta(x) := a/2 - d(x, A) > 0$  and

$$\begin{aligned} \text{dia}\{B(x, d(x, A) + \delta(x)/2) \cap A\} &\leq 2(d(x, A) + \delta(x)/2) \\ &= 2(d(x, A) + a/4 - d(x, A)/2) = d(x, A) + a/2 < a. \end{aligned}$$

**THEOREM 4.** *Let  $A = \bar{A}$  be a compact type I subset of a metric space  $(M, d)$  and let  $f: A \rightarrow M$  be a map. For each  $x \in A$  and  $\varepsilon > 0$  let  $A(f(x), \varepsilon) = \text{coS}\{B(f(x), d(f(x), A) + \varepsilon) \cap A\}$ . Then there is an  $x \in A$  such that  $x \in \bigcup_{\varepsilon > 0} A(f(x), \varepsilon)$  (this latter set will be denoted by  $A_{f(x)}$ ).*

**PROOF.** Suppose that there exists  $\delta > 0$  such that, for all  $x \in A$ ,  $x \notin A(f(x), \delta)$ . Then we take  $\delta$  in place of  $\varepsilon'(x)$  and repeat the construction of  $r$  from Proposition 4. The map  $r \circ f: A \rightarrow A$  has a fixed point [1, p. 101] which is impossible as  $(r \circ f)(x) \in A(f(x), \delta)$ . Hence there exists a sequence  $(x_n)_{n \in \mathbb{N}}$  such that  $x_n \in A(f(x_n), \delta_n)$  with  $\delta_n \rightarrow 0$ ; we may assume the sequence to converge, say to  $x$ . For any  $\delta > 0$  there exists  $n_0$  such that for every  $n \geq n_0$

$$B(f(x_n), d(f(x_n), A) + \delta_n) \subset B(f(x), d(f(x), A) + \delta).$$

Therefore  $x_n \in A(f(x), \delta)$  for  $n \geq n_0$  and  $x \in A(f(x), \delta)$ . So it must be that  $x \in A_{f(x)}$ .

Theorem 3 and Proposition 5 have locally type I analogs; the assumption that, for  $x \notin A$ ,  $d(x, A) = d(x, \partial A)$  can be omitted.

## REFERENCES

1. K. Borsuk, *Theory of retracts*, PWN, Warszawa, 1967.
2. D. W. Curtis, *Some theorems and examples on local equiconnectedness and its specializations*, *Fund. Math.* **72** (1971), 101–113.
3. J. Dugundji, *Locally equiconnected spaces and absolute neighborhood retracts*, *Fund. Math.* **57** (1965), 187–193.
4. R. H. Fox, *On fiber spaces. II*, *Bull. Amer. Math. Soc.* **49** (1943), 733–735.
5. C.J. Himmelberg, *Some theorems on equiconnected and locally equiconnected spaces*, *Trans. Amer. Math. Soc.* **115** (1965), 43–53.

SCIENCE SCHOOL OF MINING AND METALLURGY, INSTITUTE OF MATHEMATICS, KRAKÓW, AL. MICKIEWICZA 30, POLAND